



# Recombinant Goat Interleukin-4 (IL4)

<b>Product Code</b>	CSB-BP011659GO
<b>Storage</b>	Store at -20°C, for extended storage, conserve at -20°C or -80°C.
<b>Uniprot No.</b>	P79155
<b>Product Type</b>	Recombinant Protein
<b>Immunogen Species</b>	Capra hircus (Goat)
<b>Purity</b>	>85% (SDS-PAGE)
<b>Sequence</b>	HKCDIT LEEIHKMLNI LTSQKNSCME LPVADVFAAP KNATEKETFC RAGIELRRIY RNHMCLNKFL GGLDRNLSSL ASKTCSVNEA KTSTSTLRDL LERLKTIMKE KYSKC
<b>Source</b>	Baculovirus
<b>Target Names</b>	IL4
<b>Protein Names</b>	Recommended name: Interleukin-4 Short name= IL-4 Alternative name(s): B-cell stimulatory factor 1 Short name= BSF-1 Lymphocyte stimulatory factor 1
<b>Expression Region</b>	25-135
<b>Notes</b>	Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.
<b>Tag Info</b>	Tag type will be determined during the manufacturing process.
<b>Protein Length</b>	Full Length of Mature Protein
<b>Target Details</b>	This protein is a pleiotropic cytokine produced by activated T cells. This cytokine is a ligand for interleukin 4 receptor. The interleukin 4 receptor also binds to IL13, which may contribute to many overlapping functions of this cytokine and IL13. STAT6, a signal transducer and activator of transcription, has been shown to play a central role in mediating the immune regulatory signal of this cytokine. This gene, IL3, IL5, IL13, and CSF2 form a cytokine gene cluster on chromosome 5q, with this gene particularly close to IL13. This gene, IL13 and IL5 are found to be regulated coordinately by several long-range regulatory elements in an over 120 kilobase range on the chromosome. Two alternatively spliced transcript variants of this gene encoding distinct isoforms have been reported.
<b>Reconstitution</b>	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL. We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference.
<b>Shelf Life</b>	The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself. Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.