





INMT Antibody

| | |
|----------------------------|---|
| Product Code | CSB-PA011723LA01HU |
| Storage | Upon receipt, store at -20°C or -80°C. Avoid repeated freeze. |
| Uniprot No. | O95050 |
| Immunogen | Recombinant Human Indolethylamine N-methyltransferase protein (1-263AA) |
| Raised In | Rabbit |
| Species Reactivity | Human, Mouse |
| Tested Applications | ELISA, WB, IHC, IF; Recommended dilution: WB:1:1000-1:5000, IHC:1:20-1:200, IF:1:50-1:200 |
| Form | Liquid |
| Conjugate | Non-conjugated |
| Storage Buffer | Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 |
| Purification Method | >95%, Protein G purified |
| Isotype | IgG |
| Clonality | Polyclonal |
| Alias | Indolethylamine N-methyltransferase (Indolamine N-methyltransferase) (EC 2.1.1.49) (EC 2.1.1.96) (Aromatic alkylamine N-methyltransferase) (Amine N-methyltransferase) (Arylamine N-methyltransferase) (Thioether S-methyltransferase) (TEMT), INMT |
| Immunogen Species | Homo sapiens (Human) |
| Research Area | Signal Transduction |
| Target Names | INMT |


Image



Western blot
All lanes: INMT antibody at 2µg/ml + Mouse lung tissue
Secondary
Goat polyclonal to rabbit IgG at 1/15000 dilution
Predicted band size: 29 kDa
Observed band size: 29 kDa



Immunohistochemistry of paraffin-embedded human lung cancer using CSB-PA011723LA01HU at dilution of 1:100



Immunofluorescence staining of A549 cells with CSB-PA011723LA01HU at 1:150, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).