



# Recombinant Human 3-hydroxyacyl-CoA dehydrogenase type-2 (HSD17B10)

<b>Product Code</b>	CSB-EP860776HU
<b>Relevance</b>	Functions in mitochondrial tRNA maturation. Part of mitochondrial ribonuclease P, an enzyme composed of MRPP1/TRMT10C, MRPP2/HSD17B10 and MRPP3/KIAA0391, which cleaves tRNA molecules in their 5'-ends. Catalyzes the beta-oxidation at position 17 of androgens and estrogens and has 3-alpha-hydroxysteroid dehydrogenase activity with androsterone. Catalyzes the third step in the beta-oxidation of fatty acids. Carries out oxidative conversions of 7-alpha-OH and 7-beta-OH bile acids. Also exhibits 20-beta-OH and 21-OH dehydrogenase activities with C21 steroids. By interacting with intracellular amyloid-beta, it may contribute to the neuronal dysfunction associated with Alzheimer disease (AD).
<b>Abbreviation</b>	Recombinant Human HSD17B10 protein
<b>Storage</b>	The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself. Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.
<b>Uniprot No.</b>	Q99714
<b>Alias</b>	17-beta-hydroxysteroid dehydrogenase 10 (EC:1.1.1.51) ;17-beta-HSD 103-hydroxy-2-methylbutyryl-CoA dehydrogenase (EC:1.1.1.178)3-hydroxyacyl-CoA dehydrogenase type IIEndoplasmic reticulum-associated amyloid beta-peptide-binding protein;Mitochondrial ribonuclease P protein 2 ;Mitochondrial RNase P protein 2Short chain dehydrogenase/reductase family 5C member 1Short-chain type dehydrogenase/reductase XH98G2Type II HADH
<b>Product Type</b>	Recombinant Protein
<b>Immunogen Species</b>	Homo sapiens (Human)
<b>Purity</b>	Greater than 90% as determined by SDS-PAGE.
<b>Sequence</b>	AAACRSVKGLVAVITGGASGLGLATAERLVGQGASAVLLDLPNSGGEAQAKKL GNNCVFAPADVTSEKDVQTALALAKGKFGFRVDVAVNCAGIAVASKTYNLKKG QHTLTLEDFQRVLDVNLMGTFNVIRLVAGEMGQNEPDQGGQRGVIINTASVAAF EGQVGQAAYSASKGGIVGMTLPIARDLAPIGIRVMTIAPGLFGTPLLTSLEPKVC NFLASQVPFPSRLGDPAEYAHVLVQAIENPFLNGEVIRLDGAIRMQP
<b>Research Area</b>	Transcription
<b>Source</b>	E.coli
<b>Target Names</b>	HSD17B10
<b>Expression Region</b>	2-261aa
<b>Notes</b>	Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.




**Tag Info** N-terminal 6xHis-SUMO-tagged

**Mol. Weight** 42.8kDa

**Protein Length** Full Length of Mature Protein

**Image**



(Tris-Glycine gel) Discontinuous SDS-PAGE (reduced) with 5% enrichment gel and 15% separation gel.

**Reconstitution**

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL. We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference.

**Shelf Life**

The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself. Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.