

🕜 Tel: +1-301-363-4651 🛛 🖂 Email: cusabio@cusabio.com 🥃 Website: www.cusabio.com 🍙

APOH Antibody

cycles.Purification MethodAntigen Affinity PurifiedIsotypeIgGAliasapolipoprotein H (beta-2-glycoprotein I);APOH;B2G1;BG ;Product TypePolyclonal AntibodyImmunogen SpeciesHomo sapiens (Human)Target NamesAPOHTarget DetailsApolipoprotein H has been implicated in a variety of physiologic pathway including lipoprotein metabolism, coagulation, and the production of antiphospholipid autoantibodies. APOH may be a required cofactor for a phospholipid binding by the antiphospholipid autoantibodies found in ser many patients with lupus and primary antiphospholipid syndrome, but it o seem to be required for the reactivity of antiphospholipid autoantibodies		
Uniprot No.P02749ImmunogenHuman APOHRaised InRabbitSpecies ReactivityHumanTested ApplicationsELISA,WB,IHCStorage BufferPBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoid freeze cycles.Purification MethodAntigen Affinity PurifiedIsotypeIgGAliasapolipoprotein H (beta-2-glycoprotein I);APOH;B2G1;BG ;Product TypePolyclonal AntibodyImmunogen SpeciesHomo sapiens (Human)Target NamesAPOHTarget DetailsApolipoprotein H has been implicated in a variety of physiologic pathway including lipoprotein metabolism, coagulation, and the production of antiphospholipid autoantibodies. APOH may be a required cofactor for a phospholipid binding by the antiphospholipid autoantibodies found in ser many patients with lupus and primary antiphospholipid syndrome, but it o seem to be required for the reactivity of antiphospholipid autoantibodies	Product Code	CSB-PA001939GA01HU
ImmunogenHuman APOHRaised InRabbitSpecies ReactivityHumanTested ApplicationsELISA,WB,IHCStorage BufferPBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoid freeze cycles.Purification MethodAntigen Affinity PurifiedIsotypeIgGAliasapolipoprotein H (beta-2-glycoprotein I);APOH;B2G1;BG ;Product TypePolyclonal AntibodyImmunogen SpeciesHomo sapiens (Human)Target NamesAPOHTarget DetailsApolipoprotein H has been implicated in a variety of physiologic pathway including lipoprotein metabolism, coagulation, and the production of antiphospholipid autoantibodies. APOH may be a required cofactor for a phospholipid binding by the antiphospholipid autoantibodies found in see many patients with lupus and primary antiphospholipid syndrome, but it of seem to be required for the reactivity of antiphospholipid autoantibodies	Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Raised InRabbitSpecies ReactivityHumanTested ApplicationsELISA,WB,IHCStorage BufferPBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoid freeze cycles.Purification MethodAntigen Affinity PurifiedIsotypeIgGAliasapolipoprotein H (beta-2-glycoprotein I);APOH;B2G1;BG ;Product TypePolyclonal AntibodyImmunogen SpeciesHomo sapiens (Human)Target NamesAPOHTarget DetailsApolipoprotein H has been implicated in a variety of physiologic pathway including lipoprotein metabolism, coagulation, and the production of antiphospholipid autoantibodies. APOH may be a required cofactor for a phospholipid binding by the antiphospholipid syndrome, but it is seem to be required for the reactivity of antiphospholipid autoantibodies	Uniprot No.	P02749
Species ReactivityHumanTested ApplicationsELISA,WB,IHCStorage BufferPBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoid freeze cycles.Purification MethodAntigen Affinity PurifiedIsotypeIgGAliasapolipoprotein H (beta-2-glycoprotein I);APOH;B2G1;BG ;Product TypePolyclonal AntibodyImmunogen SpeciesHomo sapiens (Human)Target NamesAPOHTarget DetailsApolipoprotein H has been implicated in a variety of physiologic pathway including lipoprotein metabolism, coagulation, and the production of antiphospholipid autoantibodies. APOH may be a required cofactor for a phospholipid binding by the antiphospholipid autoantibodies found in ser many patients with lupus and primary antiphospholipid autoantibodies	Immunogen	Human APOH
Tested ApplicationsELISA,WB,IHCStorage BufferPBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoid freeze cycles.Purification MethodAntigen Affinity PurifiedIsotypeIgGAliasapolipoprotein H (beta-2-glycoprotein I);APOH;B2G1;BG ;Product TypePolyclonal AntibodyImmunogen SpeciesHomo sapiens (Human)Target NamesAPOHTarget DetailsApolipoprotein H has been implicated in a variety of physiologic pathway including lipoprotein metabolism, coagulation, and the production of antiphospholipid autoantibodies. APOH may be a required cofactor for a phospholipid binding by the antiphospholipid autoantibodies found in ser many patients with lupus and primary antiphospholipid autoantibodies	Raised In	Rabbit
Storage BufferPBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoid freeze cycles.Purification MethodAntigen Affinity PurifiedIsotypeIgGAliasapolipoprotein H (beta-2-glycoprotein I);APOH;B2G1;BG ;Product TypePolyclonal AntibodyImmunogen SpeciesHomo sapiens (Human)Target NamesAPOHTarget DetailsApolipoprotein H has been implicated in a variety of physiologic pathway including lipoprotein metabolism, coagulation, and the production of antiphospholipid autoantibodies. APOH may be a required cofactor for a phospholipid binding by the antiphospholipid autoantibodies found in seem many patients with lupus and primary antiphospholipid syndrome, but it of seem to be required for the reactivity of antiphospholipid autoantibodies	Species Reactivity	Human
cycles.Purification MethodAntigen Affinity PurifiedIsotypeIgGAliasapolipoprotein H (beta-2-glycoprotein I);APOH;B2G1;BG ;Product TypePolyclonal AntibodyImmunogen SpeciesHomo sapiens (Human)Target NamesAPOHTarget DetailsApolipoprotein H has been implicated in a variety of physiologic pathway including lipoprotein metabolism, coagulation, and the production of antiphospholipid autoantibodies. APOH may be a required cofactor for a phospholipid binding by the antiphospholipid autoantibodies found in ser many patients with lupus and primary antiphospholipid syndrome, but it open to be required for the reactivity of antiphospholipid autoantibodies	Tested Applications	ELISA,WB,IHC
IsotypeIgGAliasapolipoprotein H (beta-2-glycoprotein I);APOH;B2G1;BG ;Product TypePolyclonal AntibodyImmunogen SpeciesHomo sapiens (Human)Target NamesAPOHTarget DetailsApolipoprotein H has been implicated in a variety of physiologic pathway including lipoprotein metabolism, coagulation, and the production of antiphospholipid autoantibodies. APOH may be a required cofactor for a phospholipid binding by the antiphospholipid autoantibodies found in ser many patients with lupus and primary antiphospholipid syndrome, but it of seem to be required for the reactivity of antiphospholipid autoantibodies	Storage Buffer	PBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoid freeze / thaw cycles.
Aliasapolipoprotein H (beta-2-glycoprotein I);APOH;B2G1;BG ;Product TypePolyclonal AntibodyImmunogen SpeciesHomo sapiens (Human)Target NamesAPOHTarget DetailsApolipoprotein H has been implicated in a variety of physiologic pathway including lipoprotein metabolism, coagulation, and the production of antiphospholipid autoantibodies. APOH may be a required cofactor for a phospholipid binding by the antiphospholipid autoantibodies found in ser many patients with lupus and primary antiphospholipid syndrome, but it o seem to be required for the reactivity of antiphospholipid autoantibodies	Purification Method	Antigen Affinity Purified
Product TypePolyclonal AntibodyImmunogen SpeciesHomo sapiens (Human)Target NamesAPOHTarget DetailsApolipoprotein H has been implicated in a variety of physiologic pathway including lipoprotein metabolism, coagulation, and the production of antiphospholipid autoantibodies. APOH may be a required cofactor for a phospholipid binding by the antiphospholipid autoantibodies found in ser many patients with lupus and primary antiphospholipid syndrome, but it of seem to be required for the reactivity of antiphospholipid autoantibodies	Isotype	IgG
Immunogen SpeciesHomo sapiens (Human)Target NamesAPOHTarget DetailsApolipoprotein H has been implicated in a variety of physiologic pathway including lipoprotein metabolism, coagulation, and the production of antiphospholipid autoantibodies. APOH may be a required cofactor for a phospholipid binding by the antiphospholipid autoantibodies found in ser many patients with lupus and primary antiphospholipid syndrome, but it of seem to be required for the reactivity of antiphospholipid autoantibodies	Alias	apolipoprotein H (beta-2-glycoprotein I);APOH;B2G1;BG ;
Target NamesAPOHTarget DetailsApolipoprotein H has been implicated in a variety of physiologic pathway including lipoprotein metabolism, coagulation, and the production of antiphospholipid autoantibodies. APOH may be a required cofactor for a phospholipid binding by the antiphospholipid autoantibodies found in ser many patients with lupus and primary antiphospholipid syndrome, but it o seem to be required for the reactivity of antiphospholipid autoantibodies	Product Type	Polyclonal Antibody
Target DetailsApolipoprotein H has been implicated in a variety of physiologic pathway including lipoprotein metabolism, coagulation, and the production of antiphospholipid autoantibodies. APOH may be a required cofactor for a phospholipid binding by the antiphospholipid autoantibodies found in ser many patients with lupus and primary antiphospholipid syndrome, but it o seem to be required for the reactivity of antiphospholipid autoantibodies	Immunogen Species	Homo sapiens (Human)
including lipoprotein metabolism, coagulation, and the production of antiphospholipid autoantibodies. APOH may be a required cofactor for a phospholipid binding by the antiphospholipid autoantibodies found in ser many patients with lupus and primary antiphospholipid syndrome, but it of seem to be required for the reactivity of antiphospholipid autoantibodies	Target Names	APOH
	Target Details	antiphospholipid autoantibodies. APOH may be a required cofactor for anionic phospholipid binding by the antiphospholipid autoantibodies found in sera of many patients with lupus and primary antiphospholipid syndrome, but it does no

1