

🕜 Tel: +1-301-363-4651 🛛 🖂 Email: cusabio@cusabio.com 🥃 Website: www.cusabio.com 🍙

BID Antibody

| cycles.Purification MethodAntigen Affinity PurifiedIsotypeIgGAliasBH3 interacting domain death agonist;BID;FP497;MGC15319;MGC42355 ;Product TypePurified Rabbit Anti human PolyClonal AntibodyImmunogen SpeciesHomo sapiens (Human)Target NamesBID | | |
|---|---------------------|---|
| Uniprot No.P55957ImmunogenHuman BIDRaised InRabbitSpecies ReactivityHumanTested ApplicationsELISA,WB,IHCStorage BufferPBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoid freeze / thav cycles.Purification MethodAntigen Affinity PurifiedIsotypeIgGAliasBH3 interacting domain death agonist;BID;FP497;MGC15319;MGC42355 ;Product TypePurified Rabbit Anti human PolyClonal AntibodyImmunogen SpeciesHomo sapiens (Human)Target NamesBIDTarget DetailsThis gene encodes a death agonist that heterodimerizes with either agonist BA or antagonist BCL2. The encoded protein is a member of the BCL-2 family of cell death regulators. It is a mediator of mitochondrial damage induced by caspase-8 (CASP8); CASP8 cleaves this encoded protein, and the COOH- terminal part translocates to mitochondria where it triggers cytochrome c release. Multiple alternatively spliced transcript variants have been found, but | Product Code | CSB-PA002698GA01HU |
| ImmunogenHuman BIDRaised InRabbitSpecies ReactivityHumanTested ApplicationsELISA,WB,IHCStorage BufferPBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoid freeze / thav cycles.Purification MethodAntigen Affinity PurifiedIsotypeIgGAliasBH3 interacting domain death agonist;BID;FP497;MGC15319;MGC42355 ;Product TypePurified Rabbit Anti human PolyClonal AntibodyImmunogen SpeciesHomo sapiens (Human)Target NamesBIDTarget DetailsThis gene encodes a death agonist that heterodimerizes with either agonist BA or antagonist BCL2. The encoded protein is a member of the BCL-2 family of caspase-8 (CASP8); CASP8 cleaves this encoded protein, and the COOH- terminal part translocates to mitochondrial damage induced by caspase-8 (CASP8); CASP8 cleaves this encoded protein, and the COOH- terminal part translocates to mitochondria where it triggers cytochrome c release. Multiple alternatively spliced transcript variants have been found, but | Storage | Upon receipt, store at -20°C or -80°C. Avoid repeated freeze. |
| Raised InRabbitSpecies ReactivityHumanTested ApplicationsELISA,WB,IHCStorage BufferPBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoid freeze / thaw cycles.Purification MethodAntigen Affinity PurifiedIsotypeIgGAliasBH3 interacting domain death agonist;BID;FP497;MGC15319;MGC42355 ;Product TypePurified Rabbit Anti human PolyClonal AntibodyImmunogen SpeciesHomo sapiens (Human)Target NamesBIDTarget DetailsThis gene encodes a death agonist that heterodimerizes with either agonist BA or antagonist BCL2. The encoded protein is a member of the BCL-2 family of cell death regulators. It is a mediator of mitochondrial damage induced by caspase-8 (CASP8); CASP8 cleaves this encoded protein, and the COOH- terminal part translocates to mitochondria where it triggers cytochrome c release. Multiple alternatively spliced transcript variants have been found, but | Uniprot No. | P55957 |
| Species ReactivityHumanTested ApplicationsELISA,WB,IHCStorage BufferPBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoid freeze / thav cycles.Purification MethodAntigen Affinity PurifiedIsotypeIgGAliasBH3 interacting domain death agonist;BID;FP497;MGC15319;MGC42355 ;Product TypePurified Rabbit Anti human PolyClonal AntibodyImmunogen SpeciesHomo sapiens (Human)Target NamesBIDTarget DetailsThis gene encodes a death agonist that heterodimerizes with either agonist BA or antagonist BCL2. The encoded protein is a member of the BCL-2 family of cell death regulators. It is a mediator of mitochondrial damage induced by caspase-8 (CASP8); CASP8 cleaves this encoded protein, and the COOH- terminal part translocates to mitochondria where it triggers cytochrome c release. Multiple alternatively spliced transcript variants have been found, but | Immunogen | Human BID |
| Tested ApplicationsELISA,WB,IHCStorage BufferPBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoid freeze / thaw cycles.Purification MethodAntigen Affinity PurifiedIsotypeIgGAliasBH3 interacting domain death agonist;BID;FP497;MGC15319;MGC42355 ;Product TypePurified Rabbit Anti human PolyClonal AntibodyImmunogen SpeciesHomo sapiens (Human)Target NamesBIDTarget DetailsThis gene encodes a death agonist that heterodimerizes with either agonist BA or antagonist BCL2. The encoded protein is a member of the BCL-2 family of cell death regulators. It is a mediator of mitochondrial damage induced by caspase-8 (CASP8); CASP8 cleaves this encoded protein, and the COOH- terminal part translocates to mitochondria where it triggers cytochrome c release. Multiple alternatively spliced transcript variants have been found, but | Raised In | Rabbit |
| Storage BufferPBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoid freeze / thav cycles.Purification MethodAntigen Affinity PurifiedIsotypeIgGAliasBH3 interacting domain death agonist;BID;FP497;MGC15319;MGC42355 ;Product TypePurified Rabbit Anti human PolyClonal AntibodyImmunogen SpeciesHomo sapiens (Human)Target NamesBIDTarget DetailsThis gene encodes a death agonist that heterodimerizes with either agonist BA or antagonist BCL2. The encoded protein is a member of the BCL-2 family of cell death regulators. It is a mediator of mitochondrial damage induced by caspase-8 (CASP8); CASP8 cleaves this encoded protein, and the COOH- terminal part translocates to mitochondria where it triggers cytochrome c release. Multiple alternatively spliced transcript variants have been found, but | Species Reactivity | Human |
| cycles.Purification MethodAntigen Affinity PurifiedIsotypeIgGAliasBH3 interacting domain death agonist;BID;FP497;MGC15319;MGC42355 ;Product TypePurified Rabbit Anti human PolyClonal AntibodyImmunogen SpeciesHomo sapiens (Human)Target NamesBIDTarget DetailsThis gene encodes a death agonist that heterodimerizes with either agonist BA or antagonist BCL2. The encoded protein is a member of the BCL-2 family of cell death regulators. It is a mediator of mitochondrial damage induced by caspase-8 (CASP8); CASP8 cleaves this encoded protein, and the COOH- terminal part translocates to mitochondria where it triggers cytochrome c release. Multiple alternatively spliced transcript variants have been found, but | Tested Applications | ELISA,WB,IHC |
| IsotypeIgGAliasBH3 interacting domain death agonist;BID;FP497;MGC15319;MGC42355 ;Product TypePurified Rabbit Anti human PolyClonal AntibodyImmunogen SpeciesHomo sapiens (Human)Target NamesBIDTarget DetailsThis gene encodes a death agonist that heterodimerizes with either agonist BA or antagonist BCL2. The encoded protein is a member of the BCL-2 family of cell death regulators. It is a mediator of mitochondrial damage induced by caspase-8 (CASP8); CASP8 cleaves this encoded protein, and the COOH- terminal part translocates to mitochondria where it triggers cytochrome c release. Multiple alternatively spliced transcript variants have been found, but | Storage Buffer | PBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoid freeze / thaw cycles. |
| AliasBH3 interacting domain death agonist;BID;FP497;MGC15319;MGC42355 ;Product TypePurified Rabbit Anti human PolyClonal AntibodyImmunogen SpeciesHomo sapiens (Human)Target NamesBIDTarget DetailsThis gene encodes a death agonist that heterodimerizes with either agonist BA or antagonist BCL2. The encoded protein is a member of the BCL-2 family of cell death regulators. It is a mediator of mitochondrial damage induced by caspase-8 (CASP8); CASP8 cleaves this encoded protein, and the COOH- terminal part translocates to mitochondria where it triggers cytochrome c release. Multiple alternatively spliced transcript variants have been found, but | Purification Method | Antigen Affinity Purified |
| Product TypePurified Rabbit Anti human PolyClonal AntibodyImmunogen SpeciesHomo sapiens (Human)Target NamesBIDTarget DetailsThis gene encodes a death agonist that heterodimerizes with either agonist BA or antagonist BCL2. The encoded protein is a member of the BCL-2 family of cell death regulators. It is a mediator of mitochondrial damage induced by caspase-8 (CASP8); CASP8 cleaves this encoded protein, and the COOH- terminal part translocates to mitochondria where it triggers cytochrome c release. Multiple alternatively spliced transcript variants have been found, but | Isotype | IgG |
| Immunogen SpeciesHomo sapiens (Human)Target NamesBIDTarget DetailsThis gene encodes a death agonist that heterodimerizes with either agonist BA or antagonist BCL2. The encoded protein is a member of the BCL-2 family of cell death regulators. It is a mediator of mitochondrial damage induced by caspase-8 (CASP8); CASP8 cleaves this encoded protein, and the COOH- terminal part translocates to mitochondria where it triggers cytochrome c release. Multiple alternatively spliced transcript variants have been found, but | Alias | BH3 interacting domain death agonist;BID;FP497;MGC15319;MGC42355 ; |
| Target Names BID Target Details This gene encodes a death agonist that heterodimerizes with either agonist BA or antagonist BCL2. The encoded protein is a member of the BCL-2 family of cell death regulators. It is a mediator of mitochondrial damage induced by caspase-8 (CASP8); CASP8 cleaves this encoded protein, and the COOH-terminal part translocates to mitochondria where it triggers cytochrome c release. Multiple alternatively spliced transcript variants have been found, but | Product Type | Purified Rabbit Anti human PolyClonal Antibody |
| Target Details This gene encodes a death agonist that heterodimerizes with either agonist BA or antagonist BCL2. The encoded protein is a member of the BCL-2 family of cell death regulators. It is a mediator of mitochondrial damage induced by caspase-8 (CASP8); CASP8 cleaves this encoded protein, and the COOH-terminal part translocates to mitochondria where it triggers cytochrome c release. Multiple alternatively spliced transcript variants have been found, but | Immunogen Species | Homo sapiens (Human) |
| or antagonist BCL2. The encoded protein is a member of the BCL-2 family of cell death regulators. It is a mediator of mitochondrial damage induced by caspase-8 (CASP8); CASP8 cleaves this encoded protein, and the COOH- terminal part translocates to mitochondria where it triggers cytochrome c release. Multiple alternatively spliced transcript variants have been found, but | Target Names | BID |
| | Target Details | cell death regulators. It is a mediator of mitochondrial damage induced by caspase-8 (CASP8); CASP8 cleaves this encoded protein, and the COOH-terminal part translocates to mitochondria where it triggers cytochrome c release. Multiple alternatively spliced transcript variants have been found, but |

1