



# CAV1 Antibody

|                            |   |
|----------------------------|---|
| <b>Product Code</b>        | CSB-PA004571GA01HU  |
| <b>Storage</b>             | Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.   |
| <b>Uniprot No.</b>         | Q03135  |
| <b>Immunogen</b>           | Human CAV1  |
| <b>Raised In</b>           | Rabbit  |
| <b>Species Reactivity</b>  | Human,Mouse,Rat   |
| <b>Tested Applications</b> | ELISA,WB,IHC  |
| <b>Storage Buffer</b>      | PBS with 0.02% Sodium Azide, 50% Glycerol, pH 7.3. -20°C, Avoid freeze / thaw cycles.   |
| <b>Purification Method</b> | Antigen Affinity purified   |
| <b>Isotype</b>             | IgG   |
| <b>Alias</b>               | caveolin 1, caveolae protein, 22kDa;CAV1;BSCL3;CAV;CGL3;MSTP085;VIP21 ;   |
| <b>Product Type</b>        | Purified Rabbit Anti human PolyClonal Antibody  |
| <b>Immunogen Species</b>   | Homo sapiens (Human)  |
| <b>Target Names</b>        | CAV1  |
| <b>Target Details</b>      | <p>The scaffolding protein encoded by this gene is the main component of the caveolae plasma membranes found in most cell types. The protein links integrin subunits to the tyrosine kinase FYN, an initiating step in coupling integrins to the Ras-ERK pathway and promoting cell cycle progression. The gene is a tumor suppressor gene candidate and a negative regulator of the Ras-p42/44 MAP kinase cascade. CAV1 and CAV2 are located next to each other on chromosome 7 and express colocalizing proteins that form a stable hetero-oligomeric complex. By using alternative initiation codons in the same reading frame, two isoforms (alpha and beta) are encoded by a single transcript from this gene.</p> |
| <b>Usage</b>               | For Research Use Only. Not for use in diagnostic or therapeutic procedures.   |