



# CHEK2 Antibody

<b>Product Code</b>	CSB-PA005339GA01HU
<b>Storage</b>	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
<b>Uniprot No.</b>	O96017
<b>Immunogen</b>	Human CHEK2
<b>Raised In</b>	Rabbit
<b>Species Reactivity</b>	Human,Mouse,Rat
<b>Tested Applications</b>	ELISA,WB
<b>Storage Buffer</b>	PBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.3. -20°C, Avoid freeze / thaw cycles.
<b>Purification Method</b>	Antigen Affinity purified
<b>Isotype</b>	IgG
<b>Alias</b>	CHK2 checkpoint homolog (S. pombe);CHEK2;CDS1;CHK2;HuCds1;LFS2;PP1425;RAD53 ;
<b>Product Type</b>	Purified Rabbit Anti human PolyClonal Antibody
<b>Immunogen Species</b>	Homo sapiens (Human)
<b>Target Names</b>	CHEK2
<b>Target Details</b>	<p>In response to DNA damage and replication blocks, cell cycle progression is halted through the control of critical cell cycle regulators. This protein is a cell cycle checkpoint regulator and putative tumor suppressor. It contains a forkhead-associated protein interaction domain essential for activation in response to DNA damage and is rapidly phosphorylated in response to replication blocks and DNA damage. When activated, the encoded protein is known to inhibit CDC25C phosphatase, preventing entry into mitosis, and has been shown to stabilize the tumor suppressor protein p53, leading to cell cycle arrest in G1. In addition, this protein interacts with and phosphorylates BRCA1, allowing BRCA1 to restore survival after DNA damage. Mutations in this gene have been linked with Li-Fraumeni syndrome, a highly penetrant familial cancer phenotype usually associated with inherited mutations in TP53. Also, mutations in this gene are thought to confer a predisposition to sarcomas, breast cancer, and brain tumors. This nuclear protein is a member of the CDS1 subfamily of serine/threonine protein kinases. Three transcript variants encoding different isoforms have been found for this gene.</p>
<b>Usage</b>	For Research Use Only. Not for use in diagnostic or therapeutic procedures.