

🕜 Tel: +1-301-363-4651 🛛 🖂 Email: cusabio@cusabio.com 🥃 Website: www.cusabio.com

ICAM3 Antibody

 CSB-PA010951GA01HU Upon receipt, store at -20°C or -80°C. Avoid repeated freeze. P32942 Human ICAM3 Rabbit Human ELISA,WB,IHC PBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoid freeze / thaw cycles.
 P32942 Human ICAM3 Rabbit Human ELISA,WB,IHC PBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoid freeze / thaw
Human ICAM3 Rabbit Human ELISA,WB,IHC PBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoid freeze / thaw
Rabbit Human ELISA,WB,IHC PBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoid freeze / thaw
Human ELISA,WB,IHC PBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoid freeze / thaw
ELISA,WB,IHC PBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoid freeze / thaw
PBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoid freeze / thaw
Antigen Affinity purified
lgG
intercellular adhesion molecule 3;ICAM3;CD50;CDW50;ICAM-R ;
Purified Rabbit Anti human PolyClonal Antibody
Homo sapiens (Human)
ICAM3
This protein is a member of the intercellular adhesion molecule (ICAM) family. All ICAM proteins are type I transmembrane glycoproteins, contain 2-9 immunoglobulin-like C2-type domains, and bind to the leukocyte adhesion LFA-1 protein. This protein is constitutively and abundantly expressed by all leucocytes and may be the most important ligand for LFA-1 in the initiation of the immune response. It functions not only as an adhesion molecule, but also a a potent signalling molecule.

1