

🕜 Tel: +1-301-363-4651 🛛 🖂 Email: cusabio@cusabio.com 🥃 Website: www.cusabio.com

MID2 Antibody

cycles.Purification MethodAntigen Affinity PurifiedIsotypeIgGAliasmidline 2;MID2;FLJ37715;FLJ41813;FXY2;RNF60;TRIM1 ;Product TypePurified Rabbit Anti human PolyClonal AntibodyImmunogen SpeciesHomo sapiens (Human)Target NamesMID2Target DetailsThis protein is a member of the tripartite motif (TRIM) family. The TRIM motion includes three zinc-binding domains, a RING, a B-box type 1 and a B-box		
Uniprot No.Q9UJV3ImmunogenHuman MID2Raised InRabbitSpecies ReactivityHuman,Mouse,RatTested ApplicationsELISA,IHCStorage BufferPBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoid freeze / cycles.Purification MethodAntigen Affinity PurifiedIsotypeIgGAliasmidline 2;MID2;FLJ37715;FLJ41813;FXY2;RNF60;TRIM1 ;Product TypePurified Rabbit Anti human PolyClonal AntibodyImmunogen SpeciesHomo sapiens (Human)Target NamesMID2Target DetailsThis protein is a member of the tripartite motif (TRIM) family. The TRIM motinculdes three zinc-binding domains, a RING, a B-box type 1 and a B-box 2, and a coiled-coil region. The protein localizes to microtubular structures cytoplasm. Alternate splicing of this gene results in two transcript variants	Product Code	CSB-PA013821GA01HU
ImmunogenHuman MID2Raised InRabbitSpecies ReactivityHuman,Mouse,RatTested ApplicationsELISA,IHCStorage BufferPBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoid freeze / cycles.Purification MethodAntigen Affinity PurifiedIsotypeIgGAliasmidline 2;MID2;FLJ37715;FLJ41813;FXY2;RNF60;TRIM1 ;Product TypePurified Rabbit Anti human PolyClonal AntibodyImmunogen SpeciesHomo sapiens (Human)Target NamesMID2Target DetailsThis protein is a member of the tripartite motif (TRIM) family. The TRIM motincludes three zinc-binding domains, a RING, a B-box type 1 and a B-box 2, and a coiled-coil region. The protein localizes to microtubular structures cytoplasm. Alternate splicing of this gene results in two transcript variants	Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Raised InRabbitSpecies ReactivityHuman,Mouse,RatTested ApplicationsELISA,IHCStorage BufferPBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoid freeze / cycles.Purification MethodAntigen Affinity PurifiedIsotypeIgGAliasmidline 2;MID2;FLJ37715;FLJ41813;FXY2;RNF60;TRIM1 ;Product TypePurified Rabbit Anti human PolyClonal AntibodyImmunogen SpeciesHomo sapiens (Human)Target NamesMID2Target DetailsThis protein is a member of the tripartite motif (TRIM) family. The TRIM motioncludes three zinc-binding domains, a RING, a B-box type 1 and a B-box 2, and a coiled-coil region. The protein localizes to microtubular structures cytoplasm. Alternate splicing of this gene results in two transcript variants	Uniprot No.	Q9UJV3
Species ReactivityHuman,Mouse,RatTested ApplicationsELISA,IHCStorage BufferPBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoid freeze / cycles.Purification MethodAntigen Affinity PurifiedIsotypeIgGAliasmidline 2;MID2;FLJ37715;FLJ41813;FXY2;RNF60;TRIM1 ;Product TypePurified Rabbit Anti human PolyClonal AntibodyImmunogen SpeciesHomo sapiens (Human)Target NamesMID2Target DetailsThis protein is a member of the tripartite motif (TRIM) family. The TRIM mo includes three zinc-binding domains, a RING, a B-box type 1 and a B-box 2, and a coiled-coil region. The protein localizes to microtubular structures cytoplasm. Alternate splicing of this gene results in two transcript variants	Immunogen	Human MID2
Tested ApplicationsELISA,IHCStorage BufferPBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoid freeze / cycles.Purification MethodAntigen Affinity PurifiedIsotypeIgGAliasmidline 2;MID2;FLJ37715;FLJ41813;FXY2;RNF60;TRIM1 ;Product TypePurified Rabbit Anti human PolyClonal AntibodyImmunogen SpeciesHomo sapiens (Human)Target NamesMID2Target DetailsThis protein is a member of the tripartite motif (TRIM) family. The TRIM motincludes three zinc-binding domains, a RING, a B-box type 1 and a B-box 2, and a coiled-coil region. The protein localizes to microtubular structures cytoplasm. Alternate splicing of this gene results in two transcript variants	Raised In	Rabbit
Storage BufferPBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoid freeze / cycles.Purification MethodAntigen Affinity PurifiedIsotypeIgGAliasmidline 2;MID2;FLJ37715;FLJ41813;FXY2;RNF60;TRIM1 ;Product TypePurified Rabbit Anti human PolyClonal AntibodyImmunogen SpeciesHomo sapiens (Human)Target NamesMID2Target DetailsThis protein is a member of the tripartite motif (TRIM) family. The TRIM moti includes three zinc-binding domains, a RING, a B-box type 1 and a B-box 2, and a coiled-coil region. The protein localizes to microtubular structures cytoplasm. Alternate splicing of this gene results in two transcript variants	Species Reactivity	Human, Mouse, Rat
cycles.Purification MethodAntigen Affinity PurifiedIsotypeIgGAliasmidline 2;MID2;FLJ37715;FLJ41813;FXY2;RNF60;TRIM1 ;Product TypePurified Rabbit Anti human PolyClonal AntibodyImmunogen SpeciesHomo sapiens (Human)Target NamesMID2Target DetailsThis protein is a member of the tripartite motif (TRIM) family. The TRIM motincludes three zinc-binding domains, a RING, a B-box type 1 and a B-box 2, and a coiled-coil region. The protein localizes to microtubular structures cytoplasm. Alternate splicing of this gene results in two transcript variants	Tested Applications	ELISA,IHC
IsotypeIgGAliasmidline 2;MID2;FLJ37715;FLJ41813;FXY2;RNF60;TRIM1 ;Product TypePurified Rabbit Anti human PolyClonal AntibodyImmunogen SpeciesHomo sapiens (Human)Target NamesMID2Target DetailsThis protein is a member of the tripartite motif (TRIM) family. The TRIM motion includes three zinc-binding domains, a RING, a B-box type 1 and a B-box 2, and a coiled-coil region. The protein localizes to microtubular structures cytoplasm. Alternate splicing of this gene results in two transcript variants	Storage Buffer	PBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoid freeze / thaw cycles.
Aliasmidline 2;MID2;FLJ37715;FLJ41813;FXY2;RNF60;TRIM1 ;Product TypePurified Rabbit Anti human PolyClonal AntibodyImmunogen SpeciesHomo sapiens (Human)Target NamesMID2Target DetailsThis protein is a member of the tripartite motif (TRIM) family. The TRIM motincludes three zinc-binding domains, a RING, a B-box type 1 and a B-box 2, and a coiled-coil region. The protein localizes to microtubular structures cytoplasm. Alternate splicing of this gene results in two transcript variants	Purification Method	Antigen Affinity Purified
Product TypePurified Rabbit Anti human PolyClonal AntibodyImmunogen SpeciesHomo sapiens (Human)Target NamesMID2Target DetailsThis protein is a member of the tripartite motif (TRIM) family. The TRIM motion includes three zinc-binding domains, a RING, a B-box type 1 and a B-box 2, and a coiled-coil region. The protein localizes to microtubular structures cytoplasm. Alternate splicing of this gene results in two transcript variants	Isotype	IgG
Immunogen SpeciesHomo sapiens (Human)Target NamesMID2Target DetailsThis protein is a member of the tripartite motif (TRIM) family. The TRIM motincludes three zinc-binding domains, a RING, a B-box type 1 and a B-box 2, and a coiled-coil region. The protein localizes to microtubular structures cytoplasm. Alternate splicing of this gene results in two transcript variants	Alias	midline 2;MID2;FLJ37715;FLJ41813;FXY2;RNF60;TRIM1 ;
Target Names MID2 Target Details This protein is a member of the tripartite motif (TRIM) family. The TRIM motion includes three zinc-binding domains, a RING, a B-box type 1 and a B-box 2, and a coiled-coil region. The protein localizes to microtubular structures cytoplasm. Alternate splicing of this gene results in two transcript variants	Product Type	Purified Rabbit Anti human PolyClonal Antibody
Target DetailsThis protein is a member of the tripartite motif (TRIM) family. The TRIM motionIncludes three zinc-binding domains, a RING, a B-box type 1 and a B-box2, and a coiled-coil region. The protein localizes to microtubular structurescytoplasm. Alternate splicing of this gene results in two transcript variants	Immunogen Species	Homo sapiens (Human)
includes three zinc-binding domains, a RING, a B-box type 1 and a B-box 2, and a coiled-coil region. The protein localizes to microtubular structures cytoplasm. Alternate splicing of this gene results in two transcript variants	Target Names	MID2
	Target Details	

1