



# PAK2 Antibody

|                            |  |
|----------------------------|--|
| <b>Product Code</b>        | CSB-PA017406GA01HU   |
| <b>Storage</b>             | Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.  |
| <b>Uniprot No.</b>         | Q13177   |
| <b>Immunogen</b>           | Human PAK2   |
| <b>Raised In</b>           | Rabbit   |
| <b>Species Reactivity</b>  | Human  |
| <b>Tested Applications</b> | ELISA, WB, IHC, IF   |
| <b>Storage Buffer</b>      | PBS with 0.02% Sodium Azide, 50% Glycerol, pH 7.3. -20°C, Avoid freeze / thaw cycles.  |
| <b>Purification Method</b> | Antigen Affinity purified  |
| <b>Isotype</b>             | IgG  |
| <b>Alias</b>               | p21 protein (Cdc42/Rac)-activated kinase 2; PAK2; PAK65; PAKgamma ;  |
| <b>Product Type</b>        | Purified Rabbit Anti Human PolyClonal Antibody   |
| <b>Immunogen Species</b>   | Homo sapiens (Human)   |
| <b>Target Names</b>        | PAK2   |
| <b>Target Details</b>      | The p21 activated kinases (PAK) are critical effectors that link Rho GTPases to cytoskeleton reorganization and nuclear signaling. The PAK proteins are a family of serine/threonine kinases that serve as targets for the small GTP binding proteins, CDC42 and RAC1, and have been implicated in a wide range of biological activities. This protein is activated by proteolytic cleavage during caspase-mediated apoptosis, and may play a role in regulating the apoptotic events in the dying cell. |
| <b>Usage</b>               | For Research Use Only. Not for use in diagnostic or therapeutic procedures.  |