

defined.





PLAUR Antibody

| Product Code | CSB-PA018122GA01HU |
|----------------------------|--|
| Storage | Upon receipt, store at -20°C or -80°C. Avoid repeated freeze. |
| Uniprot No. | Q03405 |
| Immunogen | Human PLAUR |
| Raised In | Rabbit |
| Species Reactivity | Human |
| Tested Applications | ELISA,WB |
| Storage Buffer | PBS with 0.02% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoid freeze / thaw cycles. |
| Purification Method | Antigen Affinity purified |
| Isotype | IgG |
| Alias | plasminogen activator, urokinase receptor;PLAUR;CD87;UPAR;URKR; |
| Product Type | Purified Rabbit Anti human PolyClonal Antibody |
| Immunogen Species | Homo sapiens (Human) |
| Target Names | PLAUR |
| Target Details | This gene encodes the receptor for urokinase plasminogen activator and, given its role in localizing and promoting plasmin formation, likely influences many normal and pathological processes related to cell-surface plasminogen activation and localized degradation of the extracellular matrix. It binds both the proprotein and mature forms of urokinase plasminogen activator and permits the activation of the receptor-bound pro-enzyme by plasmin. The protein lacks transmembrane or cytoplasmic domains and may be anchored to the plasma membrane by a glycosyl-phosphatidylinositol (GPI) moiety following cleavage of the nascent polypeptide near its carboxy-terminus. However, a soluble protein is also produced in some cell types. Alternative splicing results in multiple transcript variants encoding different isoforms. The proprotein experiences several post-translational cleavage reactions that have not yet been fully |