

🕜 Tel: +1-301-363-4651 🛛 🖂 Email: cusabio@cusabio.com 🥃 Website: www.cusabio.com

PRPF8 Antibody

cycles.Purification MethodAntigen Affinity PurifiedIsotypeIgGAliasPRP8 pre-mRNA processing factor 8 homolog (S. cerevisiae);PRPF8;HPRP8;PRPC8;RP13 ;Product TypePurified Rabbit Anti human PolyClonal AntibodyImmunogen SpeciesHomo sapiens (Human)Target NamesPRPF8Pre-mRNA splicing occurs in 2 sequential transesterification ster is a component of both U2- and U12-dependent spliceosomes, and a set of the set of the spliceosomes, and a set of the		
Uniprot No.Q6P2Q9ImmunogenHuman PRPF8Raised InRabbitSpecies ReactivityHuman,Mouse,RatTested ApplicationsELISA,WBStorage BufferPBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.320°C, Avaccycles.Purification MethodAntigen Affinity PurifiedIsotypeIgGAliasPRP8 pre-mRNA processing factor 8 homolog (S.cerevisiae);PRPF8;HPRP8;PRPC8;RP13;Product TypePurified Rabbit Anti human PolyClonal AntibodyImmunogen SpeciesHomo sapiens (Human)Target NamesPRPF8Pre-mRNA splicing occurs in 2 sequential transesterification steris a component of both U2- and U12-dependent spliceosomes, and a set of the set of th	Code CS	SB-PA018773GA01HU
ImmunogenHuman PRPF8Raised InRabbitSpecies ReactivityHuman,Mouse,RatTested ApplicationsELISA,WBStorage BufferPBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.320°C, Avo cycles.Purification MethodAntigen Affinity PurifiedIsotypeIgGAliasPRP8 pre-mRNA processing factor 8 homolog (S. cerevisiae);PRPF8;HPRP8;PRPC8;RP13;Product TypePurified Rabbit Anti human PolyClonal AntibodyImmunogen SpeciesPRPF8Target NamesPRPF8Pre-mRNA splicing occurs in 2 sequential transesterification steries is a component of both U2- and U12-dependent spliceosomes, and a splicent of both U2- and U12-dependent splicent of both	Up	pon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Raised InRabbitSpecies ReactivityHuman,Mouse,RatTested ApplicationsELISA,WBStorage BufferPBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.320°C, Avore cycles.Purification MethodAntigen Affinity PurifiedIsotypeIgGAliasPRP8 pre-mRNA processing factor 8 homolog (S. cerevisiae);PRPF8;HPRP8;PRP63;RP13;Product TypePurified Rabbit Anti human PolyClonal AntibodyImmunogen SpeciesHomo sapiens (Human)Target NamesPRP8Pre-mRNA splicing occurs in 2 sequential transesterification steris a component of both U2- and U12-dependent spliceosomes, and a splice sp	No. Qe	6P2Q9
Species ReactivityHuman,Mouse,RatTested ApplicationsELISA,WBStorage BufferPBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.320°C, Avered cycles.Purification MethodAntigen Affinity PurifiedIsotypeIgGAliasPRP8 pre-mRNA processing factor 8 homolog (S. cerevisiae);PRPF8;HPRP8;PRP8;PRPC8;RP13;Product TypePurified Rabbit Anti human PolyClonal AntibodyImmunogen SpeciesHomo sapiens (Human)Target NamesPRPF8Target DetailsPre-mRNA splicing occurs in 2 sequential transesterification stere is a component of both U2- and U12-dependent spliceosomes, and the splice of	ygen Hւ	uman PRPF8
Tested ApplicationsELISA,WBStorage BufferPBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.320°C, Avered cycles.Purification MethodAntigen Affinity PurifiedIsotypeIgGAliasPRP8 pre-mRNA processing factor 8 homolog (S. cerevisiae);PRPF8;HPRP8;PRP8;PRPC8;RP13;Product TypePurified Rabbit Anti human PolyClonal AntibodyImmunogen SpeciesHomo sapiens (Human)Target NamesPRPF8Target DetailsPre-mRNA splicing occurs in 2 sequential transesterification steries a component of both U2- and U12-dependent spliceosomes, and a splice sequential transesterification steries a component of both U2- and U12-dependent spliceosomes, and a splicent	In Ra	abbit
Storage BufferPBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.320°C, Average BufferPurification MethodAntigen Affinity PurifiedIsotypeIgGAliasPRP8 pre-mRNA processing factor 8 homolog (S. cerevisiae);PRPF8;HPRP8;PRP68;PRPC8;RP13;Product TypePurified Rabbit Anti human PolyClonal AntibodyImmunogen SpeciesHomo sapiens (Human)Target NamesPRPF8Target DetailsPre-mRNA splicing occurs in 2 sequential transesterification steries a component of both U2- and U12-dependent spliceosomes, and a splice start of the splice s	Reactivity Hu	uman,Mouse,Rat
cycles.Purification MethodAntigen Affinity PurifiedIsotypeIgGAliasPRP8 pre-mRNA processing factor 8 homolog (S. cerevisiae);PRPF8;HPRP8;PRPC8;RP13 ;Product TypePurified Rabbit Anti human PolyClonal AntibodyImmunogen SpeciesHomo sapiens (Human)Target NamesPRPF8Pre-mRNA splicing occurs in 2 sequential transesterification ster is a component of both U2- and U12-dependent spliceosomes, and a set of the set of the spliceosomes, and a set of the	Applications EL	LISA,WB
IsotypeIgGAliasPRP8 pre-mRNA processing factor 8 homolog (S. cerevisiae);PRPF8;HPRP8;PRP8;PRPC8;RP13 ;Product TypePurified Rabbit Anti human PolyClonal AntibodyImmunogen SpeciesHomo sapiens (Human)Target NamesPRPF8Target DetailsPre-mRNA splicing occurs in 2 sequential transesterification steries is a component of both U2- and U12-dependent spliceosomes, and the second		BS with 0.1% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoid freeze / thaw /cles.
AliasPRP8 pre-mRNA processing factor 8 homolog (S. cerevisiae);PRPF8;HPRP8;PRP8;PRPC8;RP13;Product TypePurified Rabbit Anti human PolyClonal AntibodyImmunogen SpeciesHomo sapiens (Human)Target NamesPRPF8Target DetailsPre-mRNA splicing occurs in 2 sequential transesterification steris is a component of both U2- and U12-dependent spliceosomes, and the second sec	tion Method Ar	ntigen Affinity Purified
cerevisiae);PRPF8;HPRP8;PRP8;PRPC8;RP13 ;Product TypePurified Rabbit Anti human PolyClonal AntibodyImmunogen SpeciesHomo sapiens (Human)Target NamesPRPF8Target DetailsPre-mRNA splicing occurs in 2 sequential transesterification steris is a component of both U2- and U12-dependent spliceosomes, and the second sec	Ig(G
Immunogen SpeciesHomo sapiens (Human)Target NamesPRPF8Target DetailsPre-mRNA splicing occurs in 2 sequential transesterification steris a component of both U2- and U12-dependent spliceosomes, a		
Target Names PRPF8 Target Details Pre-mRNA splicing occurs in 2 sequential transesterification sterilis a component of both U2- and U12-dependent spliceosomes,	t Type Pu	urified Rabbit Anti human PolyClonal Antibody
Target Details Pre-mRNA splicing occurs in 2 sequential transesterification sterilis a component of both U2- and U12-dependent spliceosomes, and U12-d	ogen Species Ho	omo sapiens (Human)
is a component of both U2- and U12-dependent spliceosomes, a	Names PF	RPF8
several WD repeats, which function in protein-protein interaction	is : es se ha	re-mRNA splicing occurs in 2 sequential transesterification steps. This protein a component of both U2- and U12-dependent spliceosomes, and found to be ssential for the catalytic step II in pre-mRNA splicing process. It contains everal WD repeats, which function in protein-protein interactions. This protein as a sequence similarity to yeast Prp8 protein. This gene is a candidate gene ir autosomal dominant retinitis pigmentosa.

1