

🕜 Tel: +1-301-363-4651 🛛 🖂 Email: cusabio@cusabio.com 🥃 Website: www.cusabio.com

SYMPK Antibody

cycles.Purification MethodAntigen Affinity PurifiedIsotypeIgGAliassymplekin;SYMPK;FLJ27092;SPK;SYM ;Product TypePurified Rabbit Anti human PolyClonal AntibodyImmunogen SpeciesHomo sapiens (Human)Target NamesSYMPKTarget DetailsThis gene encodes a nuclear protein that functions in the regulation of polyadenylation and promotes gene expression. The protein forms a high- molecular weight complex with components of the polyadenylation machinery. If is thought to serve as a scaffold for recruiting regulatory factors to the polyadenylation complex. It also participates in 3 -end maturation of histone		
Uniprot No.Q92797UmmunogenHuman SYMPKRaised InRabbitSpecies ReactivityHuman,Mouse,RatTested ApplicationsELISA,WB,IHCStorage BufferPBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoid freeze / thaw cycles.Purification MethodAntigen Affinity PurifiedIsotypeIgGAliassymplekin;SYMPK;FLJ27092;SPK;SYM ;Product TypePurified Rabbit Anti human PolyClonal AntibodyImmunogen SpeciesHomo sapiens (Human)Target NamesSYMPKTarget DetailsThis gene encodes a nuclear protein that functions in the regulation of polyadenylation and promotes gene expression. The protein forms a high- molecular weight complex with components of the polyadenylation machinery. It is thought to serve as a scaffold for recruiting regulatory factors to the polyadenylation complex. It also participates in 3 -end maturation of histone mRNAs, which do not undergo polyadenylation. The protein also localizes to the	Product Code	CSB-PA023001GA01HU
ImmunogenHuman SYMPKRaised InRabbitSpecies ReactivityHuman,Mouse,RatTested ApplicationsELISA,WB,IHCStorage BufferPBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoid freeze / thaw cycles.Purification MethodAntigen Affinity PurifiedIsotypeIgGAliassymplekin;SYMPK;FLJ27092;SPK;SYM ;Product TypePurified Rabbit Anti human PolyClonal AntibodyImmunogen SpeciesHomo sapiens (Human)Target NamesSYMPKTarget DetailsThis gene encodes a nuclear protein that functions in the regulation of polyadenylation and promotes gene expression. The protein forms a high- molecular weight complex with components of the polyadenylation machinery. It is thought to serve as a scaffold for recruiting regulatory factors to the polyadenylation complex. It also participates in 3 -end maturation of histone mRNAs, which do not undergo polyadenylation. The protein also localizes to the	Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Raised InRabbitSpecies ReactivityHuman,Mouse,RatTested ApplicationsELISA,WB,IHCStorage BufferPBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoid freeze / thaw cycles.Purification MethodAntigen Affinity PurifiedIsotypeIgGAliassymplekin;SYMPK;FLJ27092;SPK;SYM ;Product TypePurified Rabbit Anti human PolyClonal AntibodyImmunogen SpeciesHomo sapiens (Human)Target NamesSYMPKSymPkThis gene encodes a nuclear protein that functions in the regulation of polyadenylation and promotes gene expression. The protein forms a high- molecular weight complex. It also participates in 3 -end maturation of histone mRNAs, which do not undergo polyadenylation. The protein also localizes to the	Uniprot No.	Q92797
Species ReactivityHuman,Mouse,RatTested ApplicationsELISA,WB,IHCStorage BufferPBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoid freeze / thaw cycles.Purification MethodAntigen Affinity PurifiedIsotypeIgGAliassymplekin;SYMPK;FLJ27092;SPK;SYM ;Product TypePurified Rabbit Anti human PolyClonal AntibodyImmunogen SpeciesHomo sapiens (Human)Target NamesSYMPKTarget DetailsThis gene encodes a nuclear protein that functions in the regulation of polyadenylation and promotes gene expression. The protein forms a high- molecular weight complex with components of the polyadenylation machinery. It is thought to serve as a scaffold for recruiting regulatory factors to the polyadenylation complex. It also participates in 3 -end maturation of histone mRNAs, which do not undergo polyadenylation. The protein also localizes to the	Immunogen	Human SYMPK
Tested ApplicationsELISA,WB,IHCStorage BufferPBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoid freeze / thaw cycles.Purification MethodAntigen Affinity PurifiedIsotypeIgGAliassymplekin;SYMPK;FLJ27092;SPK;SYM ;Product TypePurified Rabbit Anti human PolyClonal AntibodyImmunogen SpeciesHomo sapiens (Human)Target NamesSYMPKTarget DetailsThis gene encodes a nuclear protein that functions in the regulation of polyadenylation and promotes gene expression. The protein forms a high- molecular weight complex with components of the polyadenylation machinery. It is thought to serve as a scaffold for recruiting regulatory factors to the polyadenylation complex. It also participates in 3 -end maturation of histone mRNAs, which do not undergo polyadenylation. The protein also localizes to the	Raised In	Rabbit
Storage BufferPBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoid freeze / thaw cycles.Purification MethodAntigen Affinity PurifiedIsotypeIgGAliassymplekin;SYMPK;FLJ27092;SPK;SYM ;Product TypePurified Rabbit Anti human PolyClonal AntibodyImmunogen SpeciesHomo sapiens (Human)Target NamesSYMPKTarget DetailsThis gene encodes a nuclear protein that functions in the regulation of polyadenylation and promotes gene expression. The protein forms a high- molecular weight complex. It also participates in 3 -end maturation of histone mRNAs, which do not undergo polyadenylation. The protein also localizes to the	Species Reactivity	Human,Mouse,Rat
cycles.Purification MethodAntigen Affinity PurifiedIsotypeIgGAliassymplekin;SYMPK;FLJ27092;SPK;SYM ;Product TypePurified Rabbit Anti human PolyClonal AntibodyImmunogen SpeciesHomo sapiens (Human)Target NamesSYMPKTarget DetailsThis gene encodes a nuclear protein that functions in the regulation of polyadenylation and promotes gene expression. The protein forms a high- molecular weight complex with components of the polyadenylation machinery. It is thought to serve as a scaffold for recruiting regulatory factors to the polyadenylation complex. It also participates in 3 -end maturation of histone mRNAs, which do not undergo polyadenylation. The protein also localizes to the	Tested Applications	ELISA,WB,IHC
IsotypeIgGAliassymplekin;SYMPK;FLJ27092;SPK;SYM ;Product TypePurified Rabbit Anti human PolyClonal AntibodyImmunogen SpeciesHomo sapiens (Human)Target NamesSYMPKTarget DetailsThis gene encodes a nuclear protein that functions in the regulation of polyadenylation and promotes gene expression. The protein forms a high- molecular weight complex with components of the polyadenylation machinery. It is thought to serve as a scaffold for recruiting regulatory factors to the polyadenylation complex. It also participates in 3 -end maturation of histone mRNAs, which do not undergo polyadenylation. The protein also localizes to the	Storage Buffer	PBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoid freeze / thaw cycles.
Aliassymplekin;SYMPK;FLJ27092;SPK;SYM ;Product TypePurified Rabbit Anti human PolyClonal AntibodyImmunogen SpeciesHomo sapiens (Human)Target NamesSYMPKTarget DetailsThis gene encodes a nuclear protein that functions in the regulation of polyadenylation and promotes gene expression. The protein forms a high- molecular weight complex with components of the polyadenylation machinery. It is thought to serve as a scaffold for recruiting regulatory factors to the polyadenylation complex. It also participates in 3 -end maturation of histone mRNAs, which do not undergo polyadenylation. The protein also localizes to the	Purification Method	Antigen Affinity Purified
Product TypePurified Rabbit Anti human PolyClonal AntibodyImmunogen SpeciesHomo sapiens (Human)Target NamesSYMPKTarget DetailsThis gene encodes a nuclear protein that functions in the regulation of polyadenylation and promotes gene expression. The protein forms a high- molecular weight complex with components of the polyadenylation machinery. It is thought to serve as a scaffold for recruiting regulatory factors to the polyadenylation complex. It also participates in 3 -end maturation of histone mRNAs, which do not undergo polyadenylation. The protein also localizes to the	Isotype	IgG
Immunogen SpeciesHomo sapiens (Human)Target NamesSYMPKTarget DetailsThis gene encodes a nuclear protein that functions in the regulation of polyadenylation and promotes gene expression. The protein forms a high- molecular weight complex with components of the polyadenylation machinery. It is thought to serve as a scaffold for recruiting regulatory factors to the polyadenylation complex. It also participates in 3 -end maturation of histone mRNAs, which do not undergo polyadenylation. The protein also localizes to the	Alias	symplekin;SYMPK;FLJ27092;SPK;SYM ;
Target NamesSYMPKTarget DetailsThis gene encodes a nuclear protein that functions in the regulation of polyadenylation and promotes gene expression. The protein forms a high- molecular weight complex with components of the polyadenylation machinery. It is thought to serve as a scaffold for recruiting regulatory factors to the polyadenylation complex. It also participates in 3 -end maturation of histone mRNAs, which do not undergo polyadenylation. The protein also localizes to the	Product Type	Purified Rabbit Anti human PolyClonal Antibody
Target DetailsThis gene encodes a nuclear protein that functions in the regulation of polyadenylation and promotes gene expression. The protein forms a high- molecular weight complex with components of the polyadenylation machinery. It is thought to serve as a scaffold for recruiting regulatory factors to the polyadenylation complex. It also participates in 3 -end maturation of histone mRNAs, which do not undergo polyadenylation. The protein also localizes to the	Immunogen Species	Homo sapiens (Human)
polyadenylation and promotes gene expression. The protein forms a high- molecular weight complex with components of the polyadenylation machinery. It is thought to serve as a scaffold for recruiting regulatory factors to the polyadenylation complex. It also participates in 3 -end maturation of histone mRNAs, which do not undergo polyadenylation. The protein also localizes to the	Target Names	SYMPK
	Target Details	polyadenylation and promotes gene expression. The protein forms a high- molecular weight complex with components of the polyadenylation machinery. It is thought to serve as a scaffold for recruiting regulatory factors to the polyadenylation complex. It also participates in 3 -end maturation of histone mRNAs, which do not undergo polyadenylation. The protein also localizes to the

1