







## TAF9 Antibody

<b>Product Code</b>	CSB-PA023100GA01HU
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Uniprot No.	Q16594
Immunogen	Human TAF9
Raised In	Rabbit
Species Reactivity	Human,Mouse,Rat
Tested Applications	ELISA,WB
Storage Buffer	PBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoid freeze / thaw cycles.
<b>Purification Method</b>	Antigen Affinity Purified
Isotype	IgG
Alias	TAF9 RNA polymerase II, TATA box binding protein (TBP)-associated factor, 32kDa;TAF9;AD-004;AK6;CGI-137;CINAP;CIP;MGC1603;MGC3647;MGC5067; MGC:1603;MGC:3647;MGC:5067;TAF2G;TAFII31;TAFII32;TAFIID32;
Product Type	Purified Rabbit Anti human PolyClonal Antibody
Immunogen Species	Homo sapiens (Human)
Target Names	TAF9
Target Details	Initiation of transcription by RNA polymerase II requires the activities of more than 70 polypeptides. The protein that coordinates these activities is transcription factor IID (TFIID), which binds to the core promoter to position the polymerase properly, serves as the scaffold for assembly of the remainder of the transcription complex, and acts as a channel for regulatory signals. TFIID is composed of the TATA-binding protein (TBP) and a group of evolutionarily conserved proteins known as TBP-associated factors or TAFs. TAFs may participate in basal transcription, serve as coactivators, function in promoter recognition or modify general transcription factors (GTFs) to facilitate complex assembly and transcription initiation. This gene encodes one of the smaller subunits of TFIID that binds to the basal transcription factor GTF2B as well as to several transcriptional activators such as p53 and VP16. A similar but distinct gene (TAF9L) has been found on the X chromosome and a pseudogene has

been identified on chromosome 19. Alternative splicing results in multiple

transcript variants encoding different isoforms.