



# CCT6A Antibody

|                            |   |
|----------------------------|---|
| <b>Product Code</b>        | CSB-PA004865GA01HU  |
| <b>Storage</b>             | Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.   |
| <b>Uniprot No.</b>         | P40227  |
| <b>Immunogen</b>           | Human CCT6A-Specific  |
| <b>Raised In</b>           | Rabbit  |
| <b>Species Reactivity</b>  | Human   |
| <b>Tested Applications</b> | ELISA, WB   |
| <b>Storage Buffer</b>      | PBS with 0.02% Sodium Azide, 50% Glycerol, pH 7.3. -20°C, Avoid freeze / thaw cycles.   |
| <b>Purification Method</b> | Antigen Affinity purified   |
| <b>Isotype</b>             | IgG   |
| <b>Alias</b>               | chaperonin containing TCP1, subunit 6A (zeta 1); CCT6A; CCT-zeta; CCT-zeta-1; CCT6; Cctz; HTR3; MGC126214; MGC126215; MoDP-2; TCP-1-zeta; TCP20; TCPZ; TTCP20 ;   |
| <b>Product Type</b>        | Purified Rabbit Anti Human PolyClonal Antibody  |
| <b>Immunogen Species</b>   | Homo sapiens (Human)  |
| <b>Target Names</b>        | CCT6A   |
| <b>Target Details</b>      | This gene encodes a molecular chaperone that is member of the chaperonin containing TCP1 complex (CCT), also known as the TCP1 ring complex (TRiC). This complex consists of two identical stacked rings, each containing eight different proteins. Unfolded polypeptides enter the central cavity of the complex and are folded in an ATP-dependent manner. The complex folds various proteins, including actin and tubulin. Alternate transcriptional splice variants of this gene, encoding different isoforms, have been characterized. |
| <b>Usage</b>               | For Research Use Only. Not for use in diagnostic or therapeutic procedures.   |