

🕜 Tel: +1-301-363-4651 🛛 🖂 Email: cusabio@cusabio.com 🥃 Website: www.cusabio.com

## **PPIF** Antibody

cycles.Purification MethodAntigen Affinity purifiedIsotypeIgGAliaspeptidylprolyl isomerase F;PPIF;CYP3;Cyp-D;FLJ90798;MGC117207 ;Product TypePurified Rabbit Anti human PolyClonal AntibodyImmunogen SpeciesHomo sapiens (Human)Target NamesPPIFTarget DetailsThis protein is a member of the peptidyl-prolyl cis-trans isomerase (PPIase) family. PPIases catalyze the cis-trans isomerization of proline imidic peptide bonds in oligopeptides and accelerate the folding of proteins. This protein is potein is potein is potein is potein is protein is potein is protein is potein is potein is protein is potein is protein is protein is protein is protein is potein is protein is protein is protein is potein is protein is potein is potein is protein is potein is protein is protein is potein is protein is prote		
Uniprot No.P30405ImmunogenHuman PPIF-SpecificRaised InRabbitSpecies ReactivityHuman,Mouse,RatTested ApplicationsELISA,WB,IHC,IFStorage BufferPBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoid freeze / th cycles.Purification MethodAntigen Affinity purifiedIsotypeIgGAliaspeptidylprolyl isomerase F;PPIF;CYP3;Cyp-D;FLJ90798;MGC117207 ;Product TypePurified Rabbit Anti human PolyClonal AntibodyImmunogen SpeciesHomo sapiens (Human)Target NamesPPIFTarget DetailsThis protein is a member of the peptidyl-prolyl cis-trans isomerase (PPIase) family. PPIases catalyze the cis-trans isomerization of proline imidic peptide bonds in oligopeptides and accelerate the folding of proteins. This protein is p of the mitochondrial permeability transition pore in the inner mitochondrial membrane. Activation of this pore is thought to be involved in the induction of membrane. Activation of this pore is thought to be involved in the induction of membrane. Activation of this pore is thought to be involved in the induction of membrane. Activation of this pore is thought to be involved in the induction of membrane. Activation of this pore is thought to be involved in the induction of membrane. Activation of this pore is thought to be involved in the induction of membrane. Activation of this pore is thought to be involved in the induction of	Product Code	CSB-PA018476GA01HU
ImmunogenHuman PPIF-SpecificRaised InRabbitSpecies ReactivityHuman,Mouse,RatTested ApplicationsELISA,WB,IHC,IFStorage BufferPBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoid freeze / th cycles.Purification MethodAntigen Affinity purifiedIsotypeIgGAliaspeptidylprolyl isomerase F;PPIF;CYP3;Cyp-D;FLJ90798;MGC117207 ;Product TypePurified Rabbit Anti human PolyClonal AntibodyImmunogen SpeciesHomo sapiens (Human)Target NamesPPIFTarget DetailsThis protein is a member of the peptidyl-prolyl cis-trans isomerase (PPIase) family. PPIases catalyze the cis-trans isomerization of proline imidic peptide bonds in oligopeptides and accelerate the folding of proteins. This protein is p of the mitochondrial permeability transition pore in the inner mitochondrial membrane. Activation of this pore is thought to be involved in the induction of membrane. Activation of this pore is thought to be involved in the induction of membrane. Activation of this pore is thought to be involved in the induction of membrane. Activation of this pore is thought to be involved in the induction of membrane. Activation of this pore is thought to be involved in the induction of	Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Raised InRabbitRaised InRabbitSpecies ReactivityHuman,Mouse,RatTested ApplicationsELISA,WB,IHC,IFStorage BufferPBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoid freeze / th cycles.Purification MethodAntigen Affinity purifiedIsotypeIgGAliaspeptidylprolyl isomerase F;PPIF;CYP3;Cyp-D;FLJ90798;MGC117207 ;Product TypePurified Rabbit Anti human PolyClonal AntibodyImmunogen SpeciesHomo sapiens (Human)Target NamesPPIFTarget DetailsThis protein is a member of the peptidyl-prolyl cis-trans isomerase (PPIase) family. PPIases catalyze the cis-trans isomerization of proteins. This protein is a of the mitochondrial permeability transition pore in the inner mitochondrial membrane. Activation of this pore is thought to be involved in the induction of the mitochondrial permeability transition pore in the inner mitochondrial membrane. Activation of this pore is thought to be involved in the induction of the mitochondrial permeability transition pore in the inner mitochondrial membrane. Activation of this pore is thought to be involved in the induction of	Uniprot No.	P30405
Species ReactivityHuman,Mouse,RatTested ApplicationsELISA,WB,IHC,IFStorage BufferPBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoid freeze / th cycles.Purification MethodAntigen Affinity purifiedIsotypeIgGAliaspeptidylprolyl isomerase F;PPIF;CYP3;Cyp-D;FLJ90798;MGC117207 ;Product TypePurified Rabbit Anti human PolyClonal AntibodyImmunogen SpeciesHomo sapiens (Human)Target NamesPPIFTarget DetailsThis protein is a member of the peptidyl-prolyl cis-trans isomerase (PPIase) family. PPIases catalyze the cis-trans isomerization of proline imidic peptide bonds in oligopeptides and accelerate the folding of proteins. This protein is	Immunogen	Human PPIF-Specific
Tested ApplicationsELISA,WB,IHC,IFStorage BufferPBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoid freeze / th cycles.Purification MethodAntigen Affinity purifiedIsotypeIgGAliaspeptidylprolyl isomerase F;PPIF;CYP3;Cyp-D;FLJ90798;MGC117207 ;Product TypePurified Rabbit Anti human PolyClonal AntibodyImmunogen SpeciesHomo sapiens (Human)Target NamesPPIFTarget DetailsThis protein is a member of the peptidyl-prolyl cis-trans isomerase (PPlase) family. PPlases catalyze the cis-trans isomerization of proline imidic peptide bonds in oligopeptides and accelerate the folding of proteins. This protein is p of the mitochondrial permeability transition pore in the inner mitochondrial membrane. Activation of this pore is thought to be involved in the induction of	Raised In	Rabbit
Storage BufferPBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoid freeze / th cycles.Purification MethodAntigen Affinity purifiedIsotypeIgGAliaspeptidylprolyl isomerase F;PPIF;CYP3;Cyp-D;FLJ90798;MGC117207 ;Product TypePurified Rabbit Anti human PolyClonal AntibodyImmunogen SpeciesHomo sapiens (Human)Target NamesPPIFTarget DetailsThis protein is a member of the peptidyl-prolyl cis-trans isomerase (PPIase) family. PPIases catalyze the cis-trans isomerization of proline imidic peptide bonds in oligopeptides and accelerate the folding of proteins. This protein is p of the mitochondrial permeability transition pore in the inner mitochondrial membrane. Activation of this pore is thought to be involved in the induction of	Species Reactivity	Human,Mouse,Rat
cycles.Purification MethodAntigen Affinity purifiedIsotypeIgGAliaspeptidylprolyl isomerase F;PPIF;CYP3;Cyp-D;FLJ90798;MGC117207 ;Product TypePurified Rabbit Anti human PolyClonal AntibodyImmunogen SpeciesHomo sapiens (Human)Target NamesPPIFTarget DetailsThis protein is a member of the peptidyl-prolyl cis-trans isomerase (PPIase) family. PPIases catalyze the cis-trans isomerization of proline imidic peptide bonds in oligopeptides and accelerate the folding of proteins. This protein is p of the mitochondrial permeability transition pore in the inner mitochondrial membrane. Activation of this pore is thought to be involved in the induction of	<b>Tested Applications</b>	ELISA,WB,IHC,IF
IsotypeIgGAliaspeptidylprolyl isomerase F;PPIF;CYP3;Cyp-D;FLJ90798;MGC117207 ;Product TypePurified Rabbit Anti human PolyClonal AntibodyImmunogen SpeciesHomo sapiens (Human)Target NamesPPIFTarget DetailsThis protein is a member of the peptidyl-prolyl cis-trans isomerase (PPIase) family. PPIases catalyze the cis-trans isomerization of proline imidic peptide bonds in oligopeptides and accelerate the folding of proteins. This protein is p of the mitochondrial permeability transition pore in the inner mitochondrial membrane. Activation of this pore is thought to be involved in the induction of	Storage Buffer	PBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.320°C, Avoid freeze / thaw cycles.
Aliaspeptidylprolyl isomerase F;PPIF;CYP3;Cyp-D;FLJ90798;MGC117207 ;Product TypePurified Rabbit Anti human PolyClonal AntibodyImmunogen SpeciesHomo sapiens (Human)Target NamesPPIFTarget DetailsThis protein is a member of the peptidyl-prolyl cis-trans isomerase (PPIase) family. PPIases catalyze the cis-trans isomerization of proline imidic peptide bonds in oligopeptides and accelerate the folding of proteins. This protein is p of the mitochondrial permeability transition pore in the inner mitochondrial membrane. Activation of this pore is thought to be involved in the induction of	Purification Method	Antigen Affinity purified
Product TypePurified Rabbit Anti human PolyClonal AntibodyImmunogen SpeciesHomo sapiens (Human)Target NamesPPIFTarget DetailsThis protein is a member of the peptidyl-prolyl cis-trans isomerase (PPIase) family. PPIases catalyze the cis-trans isomerization of proline imidic peptide bonds in oligopeptides and accelerate the folding of proteins. This protein is p of the mitochondrial permeability transition pore in the inner mitochondrial membrane. Activation of this pore is thought to be involved in the induction of	Isotype	IgG
Immunogen SpeciesHomo sapiens (Human)Target NamesPPIFTarget DetailsThis protein is a member of the peptidyl-prolyl cis-trans isomerase (PPIase) family. PPIases catalyze the cis-trans isomerization of proline imidic peptide bonds in oligopeptides and accelerate the folding of proteins. This protein is p of the mitochondrial permeability transition pore in the inner mitochondrial membrane. Activation of this pore is thought to be involved in the induction of	Alias	peptidylprolyl isomerase F;PPIF;CYP3;Cyp-D;FLJ90798;MGC117207;
Target NamesPPIFTarget DetailsThis protein is a member of the peptidyl-prolyl cis-trans isomerase (PPIase) family. PPIases catalyze the cis-trans isomerization of proline imidic peptide bonds in oligopeptides and accelerate the folding of proteins. This protein is p of the mitochondrial permeability transition pore in the inner mitochondrial membrane. Activation of this pore is thought to be involved in the induction of	Product Type	Purified Rabbit Anti human PolyClonal Antibody
Target DetailsThis protein is a member of the peptidyl-prolyl cis-trans isomerase (PPIase) family. PPIases catalyze the cis-trans isomerization of proline imidic peptide bonds in oligopeptides and accelerate the folding of proteins. This protein is p of the mitochondrial permeability transition pore in the inner mitochondrial membrane. Activation of this pore is thought to be involved in the induction of	Immunogen Species	Homo sapiens (Human)
family. PPIases catalyze the cis-trans isomerization of proline imidic peptide bonds in oligopeptides and accelerate the folding of proteins. This protein is p of the mitochondrial permeability transition pore in the inner mitochondrial membrane. Activation of this pore is thought to be involved in the induction of	Target Names	PPIF
	Target Details	family. PPIases catalyze the cis-trans isomerization of proline imidic peptide bonds in oligopeptides and accelerate the folding of proteins. This protein is part of the mitochondrial permeability transition pore in the inner mitochondrial membrane. Activation of this pore is thought to be involved in the induction of

1