




# COX5A Antibody

<b>Product Code</b>	CSB-PA00414A0Rb
<b>Storage</b>	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
<b>Uniprot No.</b>	P20674
<b>Immunogen</b>	Recombinant Human Cytochrome c oxidase subunit 5A, mitochondrial protein (42-150AA)
<b>Raised In</b>	Rabbit
<b>Species Reactivity</b>	Human
<b>Tested Applications</b>	ELISA, WB, IHC, IF; Recommended dilution: WB:1:1000-1:5000, IHC:1:20-1:200, IF:1:50-1:200
<b>Form</b>	Liquid
<b>Conjugate</b>	Non-conjugated
<b>Storage Buffer</b>	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4
<b>Purification Method</b>	>95%, Protein G purified
<b>Isotype</b>	IgG
<b>Clonality</b>	Polyclonal
<b>Alias</b>	Cytochrome c oxidase subunit 5A, mitochondrial (Cytochrome c oxidase polypeptide Va), COX5A
<b>Immunogen Species</b>	Homo sapiens (Human)
<b>Research Area</b>	Metabolism
<b>Target Names</b>	COX5A

## Image



### Western blot


All lanes: COX5A antibody at 2µg/ml + 293T whole cell lysate

### Secondary


Goat polyclonal to rabbit IgG at 1/15000 dilution

Predicted band size: 17 kDa


Observed band size: 17 kDa



Immunohistochemistry of paraffin-embedded human brain tissue using CSB-PA00414A0Rb at dilution of 1:100



Immunohistochemistry of paraffin-embedded human glioma using CSB-PA00414A0Rb at dilution of 1:100



Immunofluorescence staining of HeLa cells with CSB-PA00414A0Rb at 1:166, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).

## Usage

For Research Use Only. Not for use in diagnostic or therapeutic procedures.