



Recombinant Human Replication factor C subunit 5 (RFC5)

Product Code	CSB-MP019592HU
Storage	Store at -20°C, for extended storage, conserve at -20°C or -80°C.
Uniprot No.	P40937
Product Type	Recombinant Protein
Immunogen Species	Homo sapiens (Human)
Purity	≥85% (SDS-PAGE)
Sequence	METSALKQQE QPAATKIRNL PWVEKYRPQT LNDLISHQDI LSTIQKFINE DRLPHLLLYG PPGTGKTSTI LACAKQLYKD KEFGSMVLEL NASDDRIGDI IRGPILSFAS TRTIFKKGFK LVILDEADAM TQDAQNALRR VIEKFTENTR FCLICNYLSK IIPALQSRCT RFRFGPLTPE LMVPRLEHVV EEEKVDISED GMKALVTLSS GDMRRALNIL QSTNMAFGKV TEETVYTCTG HPLKSDIANI LDWMLNQDFT TAYRNITELK TLKGLALHDI LTEIHLFVHR VDFPSSVRIH LLTKMADIEY RLSVGTNEKI QLSSLIAAFQ VTRDLIVAEA
Source	Mammalian cell
Target Names	RFC5
Protein Names	Recommended name: Replication factor C subunit 5 Alternative name(s): Activator 1 36 kDa subunit Short name= A1 36 kDa subunit Activator 1 subunit 5 Replication factor C 36 kDa subunit Short name= RF-C 36 kDa subunit Shor
Expression Region	1-340
Notes	Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.
Tag Info	Tag type will be determined during the manufacturing process.
Protein Length	Full length protein
Target Details	The elongation of primed DNA templates by DNA polymerase delta and DNA polymerase epsilon requires the accessory proteins proliferating cell nuclear antigen (PCNA) and replication factor C (RFC). RFC, also named activator 1, is a protein complex consisting of five distinct subunits of 140, 40, 38, 37, and 36 kD. This gene encodes the 36 kD subunit. This subunit can interact with the C-terminal region of PCNA. It forms a core complex with the 38 and 40 kDa subunits. The core complex possesses DNA-dependent ATPase activity, which was found to be stimulated by PCNA in an in vitro system. Alternatively spliced transcript variants encoding distinct isoforms have been reported.
Reconstitution	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL. We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference.



Shelf Life

The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself.

Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.