



# NCK2 Antibody

|                            |  |
|----------------------------|--|
| <b>Product Code</b>        | CSB-PA015531HA01HU   |
| <b>Storage</b>             | Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.  |
| <b>Uniprot No.</b>         | O43639   |
| <b>Immunogen</b>           | Recombinant Human Cytoplasmic protein NCK2 protein (1-380AA)   |
| <b>Raised In</b>           | Rabbit   |
| <b>Species Reactivity</b>  | Human, Mouse   |
| <b>Tested Applications</b> | ELISA, WB, IHC, IF; Recommended dilution: WB:1:500-1:2000, IHC:1:20-1:200, IF:1:50-1:200   |
| <b>Form</b>                | Liquid   |
| <b>Conjugate</b>           | Non-conjugated   |
| <b>Storage Buffer</b>      | Preservative: 0.03% Proclin 300<br>Constituents: 50% Glycerol, 0.01M PBS, PH 7.4   |
| <b>Purification Method</b> | >95%, Protein G purified   |
| <b>Isotype</b>             | IgG  |
| <b>Clonality</b>           | Polyclonal   |
| <b>Alias</b>               | Cytoplasmic protein NCK2 (Growth factor receptor-bound protein 4) (NCK adaptor protein 2) (Nck-2) (SH2/SH3 adaptor protein NCK-beta), NCK2, GRB4 |
| <b>Immunogen Species</b>   | Homo sapiens (Human)   |
| <b>Research Area</b>       | Cell Biology   |
| <b>Target Names</b>        | NCK2   |

## Image



### Western blot

All lanes: NCK2 antibody at 8µg/ml

Lane 1: Mouse brain tissue


Lane 2: Mouse kidney tissue

### Secondary


Goat polyclonal to rabbit IgG at 1/10000 dilution

Predicted band size: 43 kDa


Observed band size: 43 kDa



Immunohistochemistry of paraffin-embedded human breast cancer using CSB-PA015531HA01HU at dilution of 1:100



Immunohistochemistry of paraffin-embedded human skin tissue using CSB-PA015531HA01HU at dilution of 1:100



Immunofluorescent analysis of HepG2 cells using CSB-PA015531HA01HU at dilution of 1:100 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L)