





AMH Antibody

Product Code	CSB-PA12639A0Rb
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Uniprot No.	P03971
Immunogen	Recombinant Human Muellerian-inhibiting factor protein (26-560AA)
Raised In	Rabbit
Species Reactivity	Human
Tested Applications	ELISA, WB, IHC, IF; Recommended dilution: WB:1:500-1:5000, IHC:1:100-1:1500, IF:1:50-1:200
Form	Liquid
Conjugate	Non-conjugated
Storage Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4
Purification Method	>95%, Protein G purified
Isotype	IgG
Clonality	Polyclonal
Alias	Muellerian-inhibiting factor (Anti-Muellerian hormone) (AMH) (Muellerian-inhibiting substance) (MIS), AMH, MIF
Immunogen Species	Homo sapiens (Human)
Research Area	Signal Transduction
Target Names	AMH


Image




Immunohistochemistry of paraffin-embedded human ovarian cancer using CSB-PA12639A0Rb at dilution of 1:100



Immunohistochemistry of paraffin-embedded human small intestine tissue using CSB-PA12639A0Rb at dilution of 1:100



Immunofluorescent analysis of MCF-7 cells using CSB-PA12639A0Rb at dilution of 1:100 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L)



Western Blot

Positive WB detected in: A549 whole cell lysate


All lanes: AMH antibody at 2.8μg/ml

Secondary

Goat polyclonal to rabbit IgG at 1/50000 dilution

Predicted band size: 60 kDa

Observed band size: 60 kDa



IHC image of CSB-PA12639A0Rb diluted at 1:1400 and staining in paraffin-embedded human prostate tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.