

Image









FLT3LG Antibody

Product Code	CSB-PA008734LA01HU
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Uniprot No.	P49771
Immunogen	Recombinant Human Fms-related tyrosine kinase 3 ligand protein (27-184AA)
Raised In	Rabbit
Species Reactivity	Human
Tested Applications	ELISA, IHC, IF; Recommended dilution: IHC:1:500-1:1000, IF:1:200-1:500
Form	Liquid
Conjugate	Non-conjugated
Storage Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4
Storage Buffer Purification Method	
	Constituents: 50% Glycerol, 0.01M PBS, PH 7.4
Purification Method	Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 >95%, Protein G purified
Purification Method	Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 >95%, Protein G purified IgG
Purification Method Isotype Clonality	Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 >95%, Protein G purified IgG Polyclonal
Purification Method Isotype Clonality Alias	Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 >95%, Protein G purified IgG Polyclonal Fms-related tyrosine kinase 3 ligand (Flt3 ligand) (Flt3L) (SL cytokine), FLT3LG
Purification Method Isotype Clonality Alias Immunogen Species	Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 >95%, Protein G purified IgG Polyclonal Fms-related tyrosine kinase 3 ligand (Flt3 ligand) (Flt3L) (SL cytokine), FLT3LG Homo sapiens (Human)



Immunofluorescence staining of MCF-7 cells with CSB-PA008734LA01HU at 1:266, counterstained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-congugated AffiniPure Goat Anti-Rabbit IgG(H+L).



IHC image of CSB-PA008734LA01HU diluted at 1:800 and staining in paraffin-embedded human lung tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.