




# IRAK4 Antibody

<b>Product Code</b>	CSB-PA011812ESR1HU
<b>Storage</b>	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
<b>Uniprot No.</b>	Q9NWZ3
<b>Immunogen</b>	Recombinant Human Interleukin-1 receptor-associated kinase 4 protein (1-220AA)
<b>Raised In</b>	Rabbit
<b>Species Reactivity</b>	Human
<b>Tested Applications</b>	ELISA, WB, IP; Recommended dilution: WB:1:500-1:2000, IP:1:200-1:2000, IP:1:200-1:2000
<b>Form</b>	Liquid
<b>Conjugate</b>	Non-conjugated
<b>Storage Buffer</b>	PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
<b>Purification Method</b>	Antigen Affinity Purified
<b>Isotype</b>	IgG
<b>Clonality</b>	Polyclonal
<b>Alias</b>	Interleukin-1 receptor-associated kinase 4 (IRAK-4) (EC 2.7.11.1) (Renal carcinoma antigen NY-REN-64), IRAK4
<b>Immunogen Species</b>	Homo sapiens (Human)
<b>Research Area</b>	Immunology
<b>Target Names</b>	IRAK4

## Image



### Western blot


All lanes: Interleukin-1 receptor-associated kinase 4 antibody at 4μg/ml + Jurkat whole cell lysate

### Secondary

Goat polyclonal to rabbit IgG at 1/10000 dilution

Predicted band size: 52, 38 kDa

Observed band size: 52 kDa



Immunoprecipitating IRAK4 in MCF-7 whole cell lysate

Lane 1: Rabbit control IgG instead of CSB-PA011812ESR1HU in MCF-7 whole cell lysate. For western blotting, a HRP-conjugated Protein G antibody was used as the secondary antibody (1/2000)

Lane 2: CSB-PA011812ESR1HU (8μg) + MCF-7 whole cell lysate (500μg)

Lane 3: MCF-7 whole cell lysate (10μg)