




Recombinant Human Protein argonaute-2 (AGO2), partial

Product Code	CSB-YP891731HU
Relevance	<p>Required for RNA-mediated gene silencing (RNAi) by the RNA-induced silencing complex (RISC). The 'minimal RISC' appears to include AGO2 bound to a short guide RNA such as a microRNA (miRNA) or short interfering RNA (siRNA). These guide RNAs direct RISC to complementary mRNAs that are targets for RISC-mediated gene silencing. The precise mechanism of gene silencing depends on the degree of complementarity between the miRNA or siRNA and its target. Binding of RISC to a perfectly complementary mRNA generally results in silencing due to endonucleolytic cleavage of the mRNA specifically by AGO2. Binding of RISC to a partially complementary mRNA results in silencing through inhibition of translation, and this is independent of endonuclease activity. May inhibit translation initiation by binding to the 7-methylguanosine cap, thereby preventing the recruitment of the translation initiation factor eIF4-E. May also inhibit translation initiation via interaction with EIF6, which itself binds to the 60S ribosomal subunit and prevents its association with the 40S ribosomal subunit. The inhibition of translational initiation leads to the accumulation of the affected mRNA in Cytoplasmic domain processing bodies (P-bodies), where mRNA degradation may subsequently occur. In some cases RISC-mediated translational repression is also observed for miRNAs that perfectly match the 3' untranslated region (3'-UTR). Can also up-regulate the translation of specific mRNAs under certain growth conditions. Binds to the AU element of the 3'-UTR of the TNF (TNF-alpha) mRNA and up-regulates translation under conditions of serum starvation. Also required for transcriptional gene silencing (TGS), in which short RNAs known as antigenic RNAs or agRNAs direct the transcriptional repression of complementary promoter regions</p>
Abbreviation	Recombinant Human AGO2 protein, partial
Storage	The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself. Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.
Uniprot No.	Q9UKV8
Product Type	Recombinant Proteins
Immunogen Species	Homo sapiens (Human)
Purity	≥ 90% as determined by SDS-PAGE.
Sequence	<p>LVVVILPGKTPVYAEVKRVGDTVLG MATQCVQMKNVQRTPQTLSNLCLKINV KLGGVNINILLPQGRPPVFQQPVIFLGADVTHPPAGDGKKPSIAAVVGSMDAHP NRYCATVRVQQHRQEIIQDLAAMVRELLIQFYKSTRFKPTRIIFYRDGVSEGQF QQVLHHELLLAIREACIKLEKDYQPGITFIVVQKRHHTRLFCTDKNERVVGKSGNIP AGTTVDTKITHPTFEFDYLC SHAGIQGTSRPSHYHVLWDDNRFSSDELQILTYQ LCHTYVRCTRSVSIPAPAYYAHLVAFRARYHLV</p>



Research Area	Epigenetics and Nuclear Signaling
Source	Yeast
Target Names	AGO2
Protein Names	Recommended name: Protein argonaute-2 Short name= Argonaute2 Short name= hAgo2 EC= 3.1.26.n2 Alternative name(s): Eukaryotic translation initiation factor 2C 2 Short name= eIF-2C 2 Short name= eIF2C 2 PAZ Piw
Expression Region	517-818aa
Notes	Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.
Tag Info	N-terminal 6xHis-tagged
Mol. Weight	36.1kDa
Protein Length	Partial

Image


(Tris-Glycine gel) Discontinuous SDS-PAGE (reduced) with 5% enrichment gel and 15% separation gel.

Shelf Life

The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself. Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.