



CD19 Monoclonal Antibody,APC Conjugated

Product Code	CSB-MA195220
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Raised In	Mouse
Species Reactivity	Human
Tested Applications	ELISA,FC;
Relevance	<p>COC19 reacts with CD19 (B4), a 90 kDa molecule, which is expressed on approximately 5-25% of human peripheral blood lymphocytes. CD19 antigen is present on human B lymphocytes at most stages of maturation, from the earliest Ig gene rearrangement in pro-B cells to mature cell, as well as malignant B cells, but is lost on maturation to plasma cells. CD19 antibody does not react with T lymphocytes, monocytes and granulocytes. CD19 is a critical signal transduction molecule that regulates B lymphocyte development, activation and differentiation. This clone is cross reactive with non-human primate.* CD19 is a key phenotyping marker of non-T cell leukemia. 1. Nadler, LM et al. (1983) J. Immunol. 131:2442. Schlossman, SL et al., eds. (1995) Leucocyte Typing V: White Cell Differentiation Antigens, Oxford University Press, New York.3. Tedder T. et al. (1994) Immunol Today. 15:437-442</p>
Form	Phosphate-buffered solution, pH7.4, containing 0.09% sodium azide and 0.2% (w/v) BSA
Isotype	IgG1
Clonality	Monoclonal
Product Type	Monoclonal Antibody
Immunogen Species	Homo sapiens (Human)
Clone No.	7F11
Usage	For Research Use Only. Not for use in diagnostic or therapeutic procedures.