

20kDa —

🕜 Tel: +1-301-363-4651 🛛 🖾 Email: cusabio@cusabio.com 🥥 Website: www.cusabio.com 🍙

## **BAG5** Antibody


CSB-PA890743ESR1HU	
Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.	
Q9UL15	
Recombinant Human BAG family molecular chaperone regulator 5 protein (198-447AA)	
Rabbit	
Human	
ELISA, WB, IHC; Recommended dilution: WB:1:500-1:2000, IHC:1:20-1:200	
Liquid	
Non-conjugated	
PBS with 0.02% sodium azide, 50% glycerol, pH7.3.	
Antigen Affinity Purified	
IgG	
Polyclonal	
BAG family molecular chaperone regulator 5 (BAG-5) (Bcl-2-associated athanogene 5), BAG5, KIAA0873	
Homo sapiens (Human)	
Cell Biology	
BAG5	
120kDa — 90kDa — 50kDa — 35kDa — 25kDa —	Western blot All lanes: BAG5 antibody at 1µg/ml + A549 whole cell lysate Secondary Goat polyclonal to rabbit IgG at 1/10000 dilution Predicted band size: 52, 57 kDa Observed band size: 52 kDa
	Upon receipt, store at -20°C or   Q9UL15   Recombinant Human BAG fam   (198-447AA)   Rabbit   Human   ELISA, WB, IHC; Recommend   Liquid   Non-conjugated   PBS with 0.02% sodium azide.   Antigen Affinity Purified   IgG   Polyclonal   BAG family molecular chapero   athanogene 5), BAG5, KIAA08   Homo sapiens (Human)   Cell Biology   BAG5

1




## **CUSABIO TECHNOLOGY LLC**


🕧 Tel: +1-301-363-4651 🛛 🖂 Email: cusabio@cusabio.com 📀 Website: www.cusabio.com 🍙



Immunohistochemistry of paraffin-embedded human breast cancer using CSB-PA890743ESR1HU at dilution of 1:100



Immunohistochemistry of paraffin-embedded human colon tissue using CSB-PA890743ESR1HU at dilution of 1:100



Immunohistochemistry of paraffin-embedded human placenta tissue using CSB-PA890743ESR1HU at dilution of 1:100