

🕜 Tel: +1-301-363-4651 🛛 🖂 Email: cusabio@cusabio.com 🥃 Website: www.cusabio.com 🍙

TNFSF13B Antibody

SB-PA897523ESR2HU	
pon receipt, store at -20°C or	-80°C. Avoid repeated freeze.
9Y275	
ecombinant Human Tumor ne otein (76-285AA)	ecrosis factor ligand superfamily member 13B
abbit	
uman, Mouse	
_ISA, WB, IHC; Recommende	ed dilution: WB:1:500-1:2000, IHC:1:20-1:200
quid	
on-conjugated	
BS with 0.02% sodium azide,	50% glycerol, pH7.3.
ntigen Affinity Purified	
G	
olyclonal	
BLyS) (B-cell-activating factor) olecule) (TNF- and APOL-rela ntigen CD257) [Cleaved into: ⁻ ember 13b, membrane form; ⁻	perfamily member 13B (B lymphocyte stimulator (BAFF) (Dendritic cell-derived TNF-like ated leukocyte expressed ligand 1) (TALL-1) (CD Tumor necrosis factor ligand superfamily Tumor necrosis factor ligand superfamily FSF13B, BAFF BLYS TALL1 TNFSF20 ZTNF4
omo sapiens (Human)	
ell Biology	
NFSF13B	
20kDa	Western blot All lanes: Tumor necrosis factor ligand superfamily member 13B antibody at 2µg/ml + Mouse liver tissue Secondary Goat polyclonal to rabbit IgG at 1/10000 dilution Predicted band size: 32, 30, 18 kDa Observed band size: 32 kDa
3	5kDa

1





CUSABIO TECHNOLOGY LLC

🕧 Tel: +1-301-363-4651 🛛 🖂 Email: cusabio@cusabio.com 📀 Website: www.cusabio.com 🍙




Immunohistochemistry of paraffin-embedded human tonsil tissue using CSB-PA897523ESR2HU at dilution of 1:100





Immunohistochemistry of paraffin-embedded human placenta tissue using CSB-PA897523ESR2HU at dilution of 1:100



Immunohistochemistry of paraffin-embedded human epityphlon tissue using CSB-PA897523ESR2HU at dilution of 1:100