





# Phospho-CDK2 (Thr160) Antibody

<b>Product Code</b>	CSB-PA912055
<b>Storage</b>	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
<b>Uniprot No.</b>	P24941
<b>Immunogen</b>	Peptide sequence around phosphorylation site of threonine 160 (T-Y-T(p)-H-E) derived from Human CDK2.
<b>Raised In</b>	Rabbit
<b>Species Reactivity</b>	Human,Mouse,Rat
<b>Specificity</b>	The antibody detects endogenous level of CDK2 only when phosphorylated at threonine 160.
<b>Tested Applications</b>	ELISA,WB,IHC;WB:1:500-1:1000,IHC:1:50-1:100
<b>Form</b>	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg <sup>2+</sup> and Ca <sup>2+</sup> ), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
<b>Purification Method</b>	Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho specific antibodies were removed by chromatography
<b>Clonality</b>	Polyclonal
<b>Alias</b>	kinase Cdk2; p33 protein kinase;
<b>Product Type</b>	Polyclonal Antibody
<b>Immunogen Species</b>	Homo sapiens (Human)
<b>Target Names</b>	CDK2

## Image



Western blot analysis of extracts from HeLa cells untreated(lane 1) or treated with UV(lane 2) using CDK2(Phospho-Thr160) Antibody .



Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using CDK2(Phospho-Thr160) Antibody (left) or the same antibody preincubated with blocking peptide(right).

**Product Modify** Phospho-Thr160



**Usage**

For Research Use Only. Not for use in diagnostic or therapeutic procedures.