





# Phospho-BCR (Tyr177) Antibody

<b>Product Code</b>	CSB-PA899546
<b>Storage</b>	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
<b>Uniprot No.</b>	P11274
<b>Immunogen</b>	Peptide sequence around phosphorylation site of tyrosine 177 (P-F-Y(p)-V-N) derived from Human Bcr.
<b>Raised In</b>	Rabbit
<b>Species Reactivity</b>	Human,Mouse
<b>Specificity</b>	The antibody detects endogenous level of Bcr only when phosphorylated at tyrosine 177.
<b>Tested Applications</b>	ELISA,WB,IHC;WB:1:500-1:1000,IHC:1:50-1:100
<b>Form</b>	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg <sup>2+</sup> and Ca <sup>2+</sup> ), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
<b>Purification Method</b>	Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho specific antibodies were removed by chromatography usi
<b>Clonality</b>	Polyclonal
<b>Alias</b>	BCR; BCR protein; BCR1
<b>Product Type</b>	Polyclonal Antibody
<b>Immunogen Species</b>	Homo sapiens (Human)
<b>Target Names</b>	BCR

## Image



Western blot analysis of extracts from K562 cells untreated or treated with H<sub>2</sub>O<sub>2</sub> using Bcr(Phospho-Tyr177) Antibody.



Immunohistochemical analysis of paraffin-embedded human tonsil tumor tissue using Bcr(Phospho-Tyr177) Antibody(left) or the same antibody preincubated with blocking peptide(right).

<b>Product Modify</b>	Phospho-Tyr177
<b>Usage</b>	For Research Use Only. Not for use in diagnostic or therapeutic procedures.