






## Phospho-PLCG2 (Tyr1217) Antibody

<b>Product Code</b>	CSB-PA227036
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Uniprot No.	P16885
Immunogen	Peptide sequence around phosphorylation site of tyrosine 1217 (F-L-Y(p)-D-T) derived from Human PLCg2.
Raised In	Rabbit
Species Reactivity	Human,Mouse,Rat
Specificity	The antibody detects endogenous level of PLC-g2 onlywhen phosphorylated at tyrosine 1217.
<b>Tested Applications</b>	ELISA,IHC;IHC:1:50-1:100
Form	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Purification Method	Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho specific antibodies were removed by chromatogramphy usi
Clonality	Polyclonal
Alias	PLC-IV; PLC-gamma2; Phospholipase C-gamma-2
Product Type	Polyclonal Antibody
Immunogen Species	Homo sapiens (Human)
Target Names	PLCG2
Image	Immunohistachemical analysis of pareffin



Immunohistochemical analysis of paraffinembedded human breast carcinoma tissue using PLC-g2(Phospho-Tyr1217) Antibody(left) or the same antibody preincubated with blocking peptide(right).

**Product Modify** 

Phospho-Tyr1217