

🕜 Tel: +1-301-363-4651 🛛 🖂 Email: cusabio@cusabio.com 🥥 Website: www.cusabio.com 🍙

IGF1R (Ab-1165/1166) Antibody

Peptide +

from Human IGF-1R . Rabbit Human,Mouse,Rat The antibody detects endogene ELISA,WB;WB:1:500-1:1000 Supplied at 1.0mg/mL in phosp pH 7.4, 150mM NaCl, 0.02% se	163~1167/1164~1168 (T-D-Y-Y-R-K) derived ous level of total IGF-1R protein.
P08069 Peptide sequence around aa.1 from Human IGF-1R . Rabbit Human,Mouse,Rat The antibody detects endogene ELISA,WB;WB:1:500-1:1000 Supplied at 1.0mg/mL in phosp pH 7.4, 150mM NaCl, 0.02% s	163~1167/1164~1168 (T-D-Y-Y-R-K) derived ous level of total IGF-1R protein.
Peptide sequence around aa.1 from Human IGF-1R . Rabbit Human,Mouse,Rat The antibody detects endogene ELISA,WB;WB:1:500-1:1000 Supplied at 1.0mg/mL in phosp pH 7.4, 150mM NaCl, 0.02% s	ous level of total IGF-1R protein. Dhate buffered saline (without Mg2+ and Ca2+), odium azide and 50% glycerol.
from Human IGF-1R . Rabbit Human,Mouse,Rat The antibody detects endogene ELISA,WB;WB:1:500-1:1000 Supplied at 1.0mg/mL in phosp pH 7.4, 150mM NaCl, 0.02% se	ous level of total IGF-1R protein. Dhate buffered saline (without Mg2+ and Ca2+), odium azide and 50% glycerol.
Human,Mouse,Rat The antibody detects endogene ELISA,WB;WB:1:500-1:1000 Supplied at 1.0mg/mL in phosp pH 7.4, 150mM NaCl, 0.02% s	bhate buffered saline (without Mg2+ and Ca2+), odium azide and 50% glycerol.
The antibody detects endogene ELISA,WB;WB:1:500-1:1000 Supplied at 1.0mg/mL in phose pH 7.4, 150mM NaCl, 0.02% se	bhate buffered saline (without Mg2+ and Ca2+), odium azide and 50% glycerol.
ELISA,WB;WB:1:500-1:1000 Supplied at 1.0mg/mL in phosp pH 7.4, 150mM NaCl, 0.02% se	bhate buffered saline (without Mg2+ and Ca2+), odium azide and 50% glycerol.
Supplied at 1.0mg/mL in phosp pH 7.4, 150mM NaCl, 0.02% s	odium azide and 50% glycerol.
pH 7.4, 150mM NaCl, 0.02% s	odium azide and 50% glycerol.
Antibodies were produced by in	manusizing relabite with events atic pentide and KIII
Antibodies were produced by immunizing rabbits with synthetic peptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific peptide.	
Polyclonal	
Insulin-like growth factor I rece	ptor; CD221; IGF1R; kinase IGF1R;
Polyclonal Antibody	
Homo sapiens (Human)	
IGF1R	
^{kD 293} 175 IGF-1R 83-	Western blot analysis of extracts from 293 cells using IGF-1R(Ab-1165/1166) Antibody and the same antibody preincubated with blocking peptide.
	175 IGF-1R

1