


🕜 Tel: +1-301-363-4651 🛛 🖂 Email: cusabio@cusabio.com 🦲 Website: www.cusabio.com 🌘

Phospho-CCNE1 (Thr395) Antibody

StorageUpon receipt, store at -20°C or -80°C. Avoid repeated freeze.Uniprot No.P24864ImmunogenPeptide sequence around phosphorylation site of threonine 395 (L-L-T(p)-P-P)derived from Human Cyclin E1Raised InRabbitSpecies ReactivityHumanSpecificityThe antibody detects endogenous level of Cyclin E1 only when phosphorylated at threonine 395.Tested ApplicationsELISA,WB;WB:1:500-1:1000FormSupplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.Purification MethodAntibodies were produced by immunizing rabbits with synthetic phosphopeptid and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho specific antibodies were removed by chromatogramphy usiClonalityPolyclonalAliasCCNE; CCNE1;Product TypePolyclonal AntibodyImmunogen SpeciesHomo sapiens (Human)Target NamesCCNE1		
Uniprot No. P24864 Immunogen Peptide sequence around phosphorylation site of threonine 395 (L-L-T(p)-P-P)derived from Human Cyclin E1 Raised In Rabbit Species Reactivity Human Specificity The antibody detects endogenous level of Cyclin E1 only when phosphorylated at threonine 395. Tested Applications ELISA,WB;WB:1:500-1:1000 Form Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Purification Method Antibodies were produced by immunizing rabbits with synthetic phosphopeptid and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho specific antibodies were removed by chromatogramphy usi Clonality Polyclonal Alias CCNE; CCNE1; Product Type Polyclonal Antibody Immunogen Species Homo sapiens (Human) Target Names CCNE1	Product Code	CSB-PA149361
ImmunogenPeptide sequence around phosphorylation site of threonine 395 (L-L-T(p)-P-P) derived from Human Cyclin E1Raised InRabbitSpecies ReactivityHumanSpecificityThe antibody detects endogenous level of Cyclin E1 only when phosphorylated at threonine 395.Tested ApplicationsELISA,WB;WB:1:500-1:1000FormSupplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.Purification MethodAntibodies were produced by immunizing rabbits with synthetic phosphopeptid and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho specific antibodies were removed by chromatogramphy usiClonalityPolyclonalAliasCCNE; CCNE1;Product TypePolyclonal AntibodyImmunogen SpeciesHomo sapiens (Human)Target NamesCCNE1	Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
P)derived from Human Cyclin E1 Raised In Rabbit Species Reactivity Human Specificity The antibody detects endogenous level of Cyclin E1 only when phosphorylated at threonine 395. Tested Applications ELISA,WB;WB:1:500-1:1000 Form Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Purification Method Antibodies were produced by immunizing rabbits with synthetic phosphopeptid and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho specific antibodies were removed by chromatogramphy usi Clonality Polyclonal Alias CCNE; CCNE1; Product Type Polyclonal Antibody Immunogen Species Homo sapiens (Human) Target Names CCNE1	Uniprot No.	P24864
Species Reactivity Human Specificity The antibody detects endogenous level of Cyclin E1 only when phosphorylated at threonine 395. Tested Applications ELISA,WB;WB:1:500-1:1000 Form Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Purification Method Antibodies were produced by immunizing rabbits with synthetic phosphopeptid and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho specific antibodies were removed by chromatogramphy usi Clonality Polyclonal Alias CCNE; CCNE1; Product Type Polyclonal Antibody Immunogen Species Homo sapiens (Human) Target Names CCNE1	Immunogen	
Specificity The antibody detects endogenous level of Cyclin E1 only when phosphorylated at threonine 395. Tested Applications ELISA,WB;WB:1:500-1:1000 Form Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Purification Method Antibodies were produced by immunizing rabbits with synthetic phosphopeptid and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho specific antibodies were removed by chromatogramphy usi Clonality Polyclonal Alias CCNE; CCNE1; Product Type Polyclonal Antibody Immunogen Species Homo sapiens (Human) Target Names CCNE1	Raised In	Rabbit
at threonine 395. Tested Applications ELISA,WB;WB:1:500-1:1000 Form Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Purification Method Antibodies were produced by immunizing rabbits with synthetic phosphopeptid and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho specific antibodies were removed by chromatogramphy usi Clonality Polyclonal Alias CCNE; CCNE1; Product Type Polyclonal Antibody Immunogen Species Homo sapiens (Human) Target Names CCNE1	Species Reactivity	Human
FormSupplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.Purification MethodAntibodies were produced by immunizing rabbits with synthetic phosphopeptid and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho specific antibodies were removed by chromatogramphy usiClonalityPolyclonalAliasCCNE; CCNE1;Product TypePolyclonal AntibodyImmunogen SpeciesHomo sapiens (Human)Target NamesCCNE1	Specificity	The antibody detects endogenous level of Cyclin E1 only when phosphorylated at threonine 395.
pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Purification Method Antibodies were produced by immunizing rabbits with synthetic phosphopeptid and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho specific antibodies were removed by chromatogramphy usi Clonality Polyclonal Alias CCNE; CCNE1; Product Type Polyclonal Antibody Immunogen Species Homo sapiens (Human) Target Names CCNE1	Tested Applications	ELISA,WB;WB:1:500-1:1000
and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho specific antibodies were removed by chromatogramphy usiClonalityPolyclonalAliasCCNE; CCNE1;Product TypePolyclonal AntibodyImmunogen SpeciesHomo sapiens (Human)Target NamesCCNE1	Form	
Alias CCNE; CCNE1; Product Type Polyclonal Antibody Immunogen Species Homo sapiens (Human) Target Names CCNE1 Image	Purification Method	
Product Type Polyclonal Antibody Immunogen Species Homo sapiens (Human) Target Names CCNE1 Image	Clonality	Polyclonal
Immunogen Species Homo sapiens (Human) Target Names CCNE1 Image	Alias	CCNE; CCNE1;
Target Names CCNE1	Product Type	Polyclonal Antibody
Image	Immunogen Species	Homo sapiens (Human)
	Target Names	CCNE1
	Image	



Western blot analysis of extracts from 293 cells untreated or treated with hydroxyurea using Cyclin E1(phospho-Thr395) Antibody.

Product Modify

Phospho-Thr395

1