




# EDA2R Antibody

<b>Product Code</b>	CSB-PA881011ESR1HU
<b>Storage</b>	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
<b>Uniprot No.</b>	Q9HAV5
<b>Immunogen</b>	Recombinant Human Tumor necrosis factor receptor superfamily member 27 protein (1-138AA)
<b>Raised In</b>	Rabbit
<b>Species Reactivity</b>	Human
<b>Tested Applications</b>	ELISA, WB, IHC; Recommended dilution: WB:1:500-1:2000, IHC:1:20-1:200
<b>Form</b>	Liquid
<b>Conjugate</b>	Non-conjugated
<b>Storage Buffer</b>	PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
<b>Purification Method</b>	Antigen Affinity Purified
<b>Isotype</b>	IgG
<b>Clonality</b>	Polyclonal
<b>Alias</b>	Tumor necrosis factor receptor superfamily member 27 (X-linked ectodysplasin-A2 receptor) (EDA-A2 receptor), EDA2R, TNFRSF27 XEDAR
<b>Immunogen Species</b>	Homo sapiens (Human)
<b>Research Area</b>	Cell Biology
<b>Target Names</b>	EDA2R

## Image



### Western blot

All lanes: Tumor necrosis factor receptor superfamily member 27 antibody at 6µg/ml

Lane 1: HepG2 whole cell lysate


Lane 2: Hela whole cell lysate

Secondary


Goat polyclonal to rabbit IgG at 1/10000 dilution

Predicted band size: 33, 36 kDa

Observed band size: 33 kDa



Immunohistochemistry of paraffin-embedded human testis tissue using CSB-PA881011ESR1HU at dilution of 1:100



Immunohistochemistry of paraffin-embedded human kidney tissue using CSB-PA881011ESR1HU at dilution of 1:100