



# RSAD2 Antibody

<b>Product Code</b>	CSB-PA848838ESR2HU
<b>Storage</b>	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
<b>Uniprot No.</b>	Q8WXG1
<b>Immunogen</b>	Recombinant Human Radical S-adenosyl methionine domain-containing protein 2 protein (42-361AA)
<b>Raised In</b>	Rabbit
<b>Species Reactivity</b>	Human
<b>Tested Applications</b>	ELISA, WB, IHC; Recommended dilution: WB:1:200-1:1000, IHC:1:20-1:200
<b>Form</b>	Liquid
<b>Conjugate</b>	Non-conjugated
<b>Storage Buffer</b>	PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
<b>Purification Method</b>	Antigen Affinity Purified
<b>Isotype</b>	IgG
<b>Clonality</b>	Polyclonal
<b>Alias</b>	Radical S-adenosyl methionine domain-containing protein 2 (Cytomegalovirus-induced gene 5 protein) (Viperin) (Virus inhibitory protein, endoplasmic reticulum-associated, interferon-inducible), RSAD2, CIG5
<b>Immunogen Species</b>	Homo sapiens (Human)
<b>Research Area</b>	Others
<b>Target Names</b>	RSAD2

## Image



Western blot


All lanes: RSAD2 antibody at 8μg/ml + Jurkat whole cell lysate

Secondary


Goat polyclonal to rabbit IgG at 1/10000 dilution

Predicted band size: 42 kDa


Observed band size: 42 kDa



Immunohistochemistry of paraffin-embedded human skin tissue using CSB-PA848838ESR2HU at dilution of 1:100



Immunohistochemistry of paraffin-embedded human colon tissue using CSB-PA848838ESR2HU at dilution of 1:100



Immunohistochemistry of paraffin-embedded human kidney tissue using CSB-PA848838ESR2HU at dilution of 1:100