





STRAP Antibody

Upon receipt, store at -20°C or -80°C. Avoid repeated freeze. Uniprot No. Q9Y3F4 Immunogen Recombinant Human Serine-threonine kinase receptor-associated protein (1-350AA) Raised In Rabbit Species Reactivity Human, Mouse Tested Applications ELISA, WB, IHC; Recommended dilution: WB:1:500-1:2000, IHC:1:20-1:200 Form Liquid Conjugate Non-conjugated Storage Buffer PBS with 0.02% sodium azide, 50% glycerol, pH7.3. Purification Method Antigen Affinity Purified Isotype IgG Clonality Polyclonal Alias Serine-threonine kinase receptor-associated protein (MAP activator with WD repeats) (UNR-interacting protein) (WD-40 repeat protein PT-WD), STRAP, MAWD UNRIP Immunogen Species Homo sapiens (Human) Research Area Epigenetics and Nuclear Signalling Target Names STRAP		
Uniprot No. Q9Y3F4 Immunogen Recombinant Human Serine-threonine kinase receptor-associated protein (1-350AA) Raised In Rabbit Species Reactivity Human, Mouse Tested Applications ELISA, WB, IHC; Recommended dilution: WB:1:500-1:2000, IHC:1:20-1:200 Form Liquid Conjugate Non-conjugated Storage Buffer PBS with 0.02% sodium azide, 50% glycerol, pH7.3. Purification Method Antigen Affinity Purified Isotype IgG Clonality Polyclonal Alias Serine-threonine kinase receptor-associated protein (MAP activator with WD repeats) (UNR-interacting protein) (WD-40 repeat protein PT-WD), STRAP, MAWD UNRIP Immunogen Species Homo sapiens (Human) Research Area Epigenetics and Nuclear Signaling Target Names STRAP	Product Code	CSB-PA896713ESR2HU
Immunogen Recombinant Human Serine-threonine kinase receptor-associated protein (1-350AA) Raised In Rabbit Species Reactivity Human, Mouse Tested Applications ELISA, WB, IHC; Recommended dilution: WB:1:500-1:2000, IHC:1:20-1:200 Form Liquid Conjugate Non-conjugated Storage Buffer PBS with 0.02% sodium azide, 50% glycerol, pH7.3. Purification Method Antigen Affinity Purified Isotype IgG Clonality Polyclonal Alias Serine-threonine kinase receptor-associated protein (MAP activator with WD repeats) (UNR-interacting protein) (WD-40 repeat protein PT-WD), STRAP, MAWD UNRIP Immunogen Species Homo sapiens (Human) Research Area Epigenetics and Nuclear Signaling Target Names Image	Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
(1-350AA) Raised In Rabbit Species Reactivity Human, Mouse Tested Applications ELISA, WB, IHC; Recommended dilution: WB:1:500-1:2000, IHC:1:20-1:200 Form Liquid Conjugate Non-conjugated Storage Buffer PBS with 0.02% sodium azide, 50% glycerol, pH7.3. Purification Method Antigen Affinity Purified Isotype IgG Clonality Polyclonal Alias Serine-threonine kinase receptor-associated protein (MAP activator with WD repeats) (UNR-interacting protein) (WD-40 repeat protein PT-WD), STRAP, MAWD UNRIP Immunogen Species Homo sapiens (Human) Research Area Epigenetics and Nuclear Signaling Target Names STRAP	Uniprot No.	Q9Y3F4
Tested Applications ELISA, WB, IHC; Recommended dilution: WB:1:500-1:2000, IHC:1:20-1:200 Form Liquid Conjugate Non-conjugated Storage Buffer PBS with 0.02% sodium azide, 50% glycerol, pH7.3. Purification Method Isotype IgG Clonality Polyclonal Alias Serine-threonine kinase receptor-associated protein (MAP activator with WD repeats) (UNR-interacting protein) (WD-40 repeat protein PT-WD), STRAP, MAWD UNRIP Immunogen Species Homo sapiens (Human) Research Area Epigenetics and Nuclear Signaling Target Names STRAP	Immunogen	·
Tested Applications	Raised In	Rabbit
Form Liquid Conjugate Non-conjugated Storage Buffer PBS with 0.02% sodium azide, 50% glycerol, pH7.3. Purification Method Antigen Affinity Purified Isotype IgG Clonality Polyclonal Alias Serine-threonine kinase receptor-associated protein (MAP activator with WD repeats) (UNR-interacting protein) (WD-40 repeat protein PT-WD), STRAP, MAWD UNRIP Immunogen Species Homo sapiens (Human) Research Area Epigenetics and Nuclear Signaling Target Names STRAP Image	Species Reactivity	Human, Mouse
Conjugate Non-conjugated PBS with 0.02% sodium azide, 50% glycerol, pH7.3. Purification Method Antigen Affinity Purified Isotype IgG Clonality Polyclonal Serine-threonine kinase receptor-associated protein (MAP activator with WD repeats) (UNR-interacting protein) (WD-40 repeat protein PT-WD), STRAP, MAWD UNRIP Immunogen Species Homo sapiens (Human) Research Area Epigenetics and Nuclear Signaling Target Names STRAP Image	Tested Applications	ELISA, WB, IHC; Recommended dilution: WB:1:500-1:2000, IHC:1:20-1:200
Storage Buffer PBS with 0.02% sodium azide, 50% glycerol, pH7.3. Purification Method Antigen Affinity Purified Isotype IgG Clonality Polyclonal Serine-threonine kinase receptor-associated protein (MAP activator with WD repeats) (UNR-interacting protein) (WD-40 repeat protein PT-WD), STRAP, MAWD UNRIP Immunogen Species Homo sapiens (Human) Research Area Epigenetics and Nuclear Signaling Target Names STRAP Image	Form	Liquid
Purification Method Antigen Affinity Purified Isotype IgG Clonality Polyclonal Alias Serine-threonine kinase receptor-associated protein (MAP activator with WD repeats) (UNR-interacting protein) (WD-40 repeat protein PT-WD), STRAP, MAWD UNRIP Immunogen Species Homo sapiens (Human) Research Area Epigenetics and Nuclear Signaling Target Names STRAP Image	Conjugate	Non-conjugated
Isotype IgG Clonality Polyclonal Alias Serine-threonine kinase receptor-associated protein (MAP activator with WD repeats) (UNR-interacting protein) (WD-40 repeat protein PT-WD), STRAP, MAWD UNRIP Immunogen Species Homo sapiens (Human) Research Area Epigenetics and Nuclear Signaling Target Names STRAP Image	Storage Buffer	PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
Clonality Polyclonal Alias Serine-threonine kinase receptor-associated protein (MAP activator with WD repeats) (UNR-interacting protein) (WD-40 repeat protein PT-WD), STRAP, MAWD UNRIP Immunogen Species Homo sapiens (Human) Research Area Epigenetics and Nuclear Signaling Target Names STRAP Image	Purification Method	Antigen Affinity Purified
Alias Serine-threonine kinase receptor-associated protein (MAP activator with WD repeats) (UNR-interacting protein) (WD-40 repeat protein PT-WD), STRAP, MAWD UNRIP Immunogen Species Homo sapiens (Human) Epigenetics and Nuclear Signaling Target Names STRAP Image	Isotype	IgG
repeats) (UNR-interacting protein) (WD-40 repeat protein PT-WD), STRAP, MAWD UNRIP Immunogen Species Homo sapiens (Human) Research Area Epigenetics and Nuclear Signaling Target Names STRAP Image	Clonality	Polyclonal
Research Area Epigenetics and Nuclear Signaling Target Names STRAP Image	Alias	repeats) (UNR-interacting protein) (WD-40 repeat protein PT-WD), STRAP,
Target Names STRAP	Immunogen Species	Homo sapiens (Human)
Image	Research Area	Epigenetics and Nuclear Signaling
Image Western blot	Target Names	STRAP
	Image	Western blot



Western blot

All lanes: STRAP antibody at 8μg/ml Lane 1: HepG2 whole cell lysate Lane 2: Hela whole cell lysate

Lane 3: Mouse brain tissue

Goat polyclonal to rabbit IgG at 1/10000 dilution

Predicted band size: 39, 40 kDa Observed band size: 39 kDa




CUSABIO TECHNOLOGY LLC












Immunohistochemistry of paraffin-embedded human colon cancer using CSB-PA896713ESR2HU at dilution of 1:100



Immunohistochemistry of paraffin-embedded human placenta tissue using CSB-PA896713ESR2HU at dilution of 1:100