




# NAA20 Antibody

<b>Product Code</b>	CSB-PA015472ESR2HU
<b>Storage</b>	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
<b>Uniprot No.</b>	P61599
<b>Immunogen</b>	Recombinant Human N-alpha-acetyltransferase 20 protein (49-178AA)
<b>Raised In</b>	Rabbit
<b>Species Reactivity</b>	Human, Rat
<b>Tested Applications</b>	ELISA, WB, IP; Recommended dilution: WB:1:1000-1:5000, IP:1:200-1:2000
<b>Form</b>	Liquid
<b>Conjugate</b>	Non-conjugated
<b>Storage Buffer</b>	PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
<b>Purification Method</b>	Antigen Affinity Purified
<b>Isotype</b>	IgG
<b>Clonality</b>	Polyclonal
<b>Alias</b>	N-alpha-acetyltransferase 20 (EC 2.3.1.254) (Methionine N-acetyltransferase) (N-acetyltransferase 5) (N-terminal acetyltransferase B complex catalytic subunit NAA20) (N-terminal acetyltransferase B complex catalytic subunit NAT5) (NatB complex subunit NAT5) (NatB catalytic subunit), NAA20, NAT5
<b>Immunogen Species</b>	Homo sapiens (Human)
<b>Research Area</b>	Others
<b>Target Names</b>	NAA20

## Image



### Western blot

All lanes: NAA20 antibody at 10μg/ml

Lane 1: Hela whole cell lysate

Lane 2: Rat brain tissue


Lane 3: 293T whole cell lysate

Secondary

Goat polyclonal to rabbit IgG at 1/10000 dilution

Predicted band size: 21, 13 kDa

Observed band size: 21 kDa



Immunoprecipitating NAA20 in HeLa whole cell lysate

Lane 1: Rabbit control IgG instead of (1 $\mu$ g) instead of CSB-PA015472ESR2HU in HeLa whole cell lysate. For western blotting, a HRP-conjugated anti-rabbit IgG, specific to the non-reduced form of IgG was used as the Secondary antibody (1/50000)

Lane 2: CSB-PA015472ESR2HU (4 $\mu$ g) + HeLa whole cell lysate (500 $\mu$ g)

Lane 3: HeLa whole cell lysate (20 $\mu$ g)