




NAA20 Antibody

Product Code	CSB-PA015472ESR2HU
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Uniprot No.	P61599
Immunogen	Recombinant Human N-alpha-acetyltransferase 20 protein (49-178AA)
Raised In	Rabbit
Species Reactivity	Human, Rat
Tested Applications	ELISA, WB, IP; Recommended dilution: WB:1:1000-1:5000, IP:1:200-1:2000
Form	Liquid
Conjugate	Non-conjugated
Storage Buffer	PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
Purification Method	Antigen Affinity Purified
Isotype	IgG
Clonality	Polyclonal
Alias	N-alpha-acetyltransferase 20 (EC 2.3.1.254) (Methionine N-acetyltransferase) (N-acetyltransferase 5) (N-terminal acetyltransferase B complex catalytic subunit NAA20) (N-terminal acetyltransferase B complex catalytic subunit NAT5) (NatB complex subunit NAT5) (NatB catalytic subunit), NAA20, NAT5
Immunogen Species	Homo sapiens (Human)
Research Area	Others
Target Names	NAA20

Image



Western blot

All lanes: NAA20 antibody at 10µg/ml

Lane 1: HeLa whole cell lysate

Lane 2: Rat brain tissue


Lane 3: 293T whole cell lysate

Secondary

Goat polyclonal to rabbit IgG at 1/10000 dilution

Predicted band size: 21, 13 kDa

Observed band size: 21 kDa



Immunoprecipitating NAA20 in HeLa whole cell lysate

Lane 1: Rabbit control IgG instead of (1 μ g) instead of CSB-PA015472ESR2HU in HeLa whole cell lysate. For western blotting, a HRP-conjugated anti-rabbit IgG, specific to the non-reduced form of IgG was used as the Secondary antibody (1/50000)

Lane 2: CSB-PA015472ESR2HU (4 μ g) + HeLa whole cell lysate (500 μ g)

Lane 3: HeLa whole cell lysate (20 μ g)

Usage

For Research Use Only. Not for use in diagnostic or therapeutic procedures.