






## MARK1/MARK2/MARK3/MARK4 Antibody

Product Code         CSB-PA003193           Storage         Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.           Uniprot No.         Q9P0L2/Q7KZI7/P27448/Q96L34           Immunogen         Synthesized peptide derived from Human MARK1/2/3/4 around the non-phosphorylation site of T215.           Raised In         Rabbit           Species Reactivity         Human,Mouse,Rat,Monkey           Tested Applications         WB, IHC, IF, ELISA; WB:1:500-1:2000, IHC:1:100-1:300, IF:1:200-1:1000, ELISA:1:10000           Form         Liquid           Conjugate         Non-conjugated           Storage Buffer         Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.           Purification Method         The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.           Isotype         IgG           Alias         MARK1; KIAA1477; MARK; Serine/threonine-protein kinase MARK1; MAP/microtubule affinity-regulating kinase 1; PAR1 homolog c; Par-1c; Par1c; MARK2; EMK1; Serine/threonine-protein kinase MARK2; ELKL motif kinase 1; EMK-1; MAP/microtubule affin           Immunogen Species         Homo sapiens (Human)           Target Names         MARK1/MARK2/MARK3/MARK4           Image		
Uniprot No. Q9P0L2/Q7KZI7/P27448/Q96L34  Immunogen Synthesized peptide derived from Human MARK1/2/3/4 around the non-phosphorylation site of T215.  Raised In Rabbit  Species Reactivity Human,Mouse,Rat,Monkey  Tested Applications WB, IHC, IF, ELISA; WB:1:500-1:2000, IHC:1:100-1:300, IF:1:200-1:1000, ELISA:1:10000  Form Liquid  Conjugate Non-conjugated  Storage Buffer Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  Purification Method The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  Isotype IgG  Alias MARK1; KIAA1477; MARK; Serine/threonine-protein kinase MARK1; MAP/microtubule affinity-regulating kinase 1; PAR1 homolog c; Par-1c; Par1c; MARK2; EMK1; Serine/threonine-protein kinase MARK2; ELKL motif kinase 1; EMK-1; MAP/microtubule affin  Immunogen Species Homo sapiens (Human)  Target Names MARK1/MARK2/MARK3/MARK4	<b>Product Code</b>	CSB-PA003193
Immunogen       Synthesized peptide derived from Human MARK1/2/3/4 around the non-phosphorylation site of T215.         Raised In       Rabbit         Species Reactivity       Human,Mouse,Rat,Monkey         Tested Applications       WB, IHC, IF, ELISA; WB:1:500-1:2000, IHC:1:100-1:300, IF:1:200-1:1000, ELISA:1:10000         Form       Liquid         Conjugate       Non-conjugated         Storage Buffer       Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.         Purification Method       The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.         Isotype       IgG         Alias       MARK1; KIAA1477; MARK; Serine/threonine-protein kinase MARK1; MAP/microtubule affinity-regulating kinase 1; PAR1 homolog c; Par-1c; Par1c; MARK2; EMK1; Serine/threonine-protein kinase MARK2; ELKL motif kinase 1; EMK-1; MAP/microtubule affin         Immunogen Species       Homo sapiens (Human)         Target Names       MARK1/MARK2/MARK3/MARK4         Image	Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
phosphorylation site of T215.  Raised In Rabbit  Species Reactivity Human,Mouse,Rat,Monkey  Tested Applications WB, IHC, IF, ELISA; WB:1:500-1:2000, IHC:1:100-1:300, IF:1:200-1:1000, ELISA:1:10000  Form Liquid  Conjugate Non-conjugated  Storage Buffer Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  Purification Method The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  Isotype IgG  Alias MARK1; KIAA1477; MARK; Serine/threonine-protein kinase MARK1; MAP/microtubule affinity-regulating kinase 1; PAR1 homolog c; Par-1c; Par1c; MARK2; EMK1; Serine/threonine-protein kinase MARK2; ELKL motif kinase 1; EMK-1; MAP/microtubule affin  Immunogen Species Homo sapiens (Human)  Target Names MARK1/MARK2/MARK3/MARK4  Image	Uniprot No.	Q9P0L2/Q7KZI7/P27448/Q96L34
Species Reactivity Human,Mouse,Rat,Monkey  Tested Applications WB, IHC, IF, ELISA; WB:1:500-1:2000, IHC:1:100-1:300, IF:1:200-1:1000, ELISA:1:10000  Form Liquid  Conjugate Non-conjugated  Storage Buffer Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  Purification Method The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  Isotype IgG  Alias MARK1; KIAA1477; MARK; Serine/threonine-protein kinase MARK1; MAP/microtubule affinity-regulating kinase 1; PAR1 homolog c; Par-1c; Par1c; MARK2; EMK1; Serine/threonine-protein kinase MARK2; ELKL motif kinase 1; EMK-1; MAP/microtubule affin	Immunogen	
Tested Applications  WB, IHC, IF, ELISA; WB:1:500-1:2000, IHC:1:100-1:300, IF:1:200-1:1000, ELISA:1:10000  Form  Liquid  Conjugate  Non-conjugated  Storage Buffer  Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  Purification Method  The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  Isotype  IgG  Alias  MARK1; KIAA1477; MARK; Serine/threonine-protein kinase MARK1; MAP/microtubule affinity-regulating kinase 1; PAR1 homolog c; Par-1c; Par1c; MARK2; EMK1; Serine/threonine-protein kinase MARK2; ELKL motif kinase 1; EMK-1; MAP/microtubule affin  Immunogen Species  Homo sapiens (Human)  Target Names  MARK1/MARK2/MARK3/MARK4	Raised In	Rabbit
ELISA:1:10000  Form Liquid  Conjugate Non-conjugated  Storage Buffer Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  Purification Method The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  Isotype IgG  Alias MARK1; KIAA1477; MARK; Serine/threonine-protein kinase MARK1; MAP/microtubule affinity-regulating kinase 1; PAR1 homolog c; Par-1c; Par1c; MARK2; EMK1; Serine/threonine-protein kinase MARK2; ELKL motif kinase 1; EMK-1; MAP/microtubule affin  Immunogen Species Homo sapiens (Human)  Target Names MARK1/MARK2/MARK3/MARK4	Species Reactivity	Human, Mouse, Rat, Monkey
Conjugate  Non-conjugated  Storage Buffer  Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  Purification Method  The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  Isotype  IgG  Alias  MARK1; KIAA1477; MARK; Serine/threonine-protein kinase MARK1; MAP/microtubule affinity-regulating kinase 1; PAR1 homolog c; Par-1c; Par1c; MARK2; EMK1; Serine/threonine-protein kinase MARK2; ELKL motif kinase 1; EMK-1; MAP/microtubule affin  Immunogen Species  Homo sapiens (Human)  Target Names  MARK1/MARK2/MARK3/MARK4	Tested Applications	
Storage Buffer  Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  Purification Method  The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  Isotype  IgG  Alias  MARK1; KIAA1477; MARK; Serine/threonine-protein kinase MARK1; MAP/microtubule affinity-regulating kinase 1; PAR1 homolog c; Par-1c; Par1c; MARK2; EMK1; Serine/threonine-protein kinase MARK2; ELKL motif kinase 1; EMK-1; MAP/microtubule affin  Immunogen Species  Homo sapiens (Human)  Target Names  MARK1/MARK2/MARK3/MARK4	Form	Liquid
Purification Method The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  Isotype IgG MARK1; KIAA1477; MARK; Serine/threonine-protein kinase MARK1; MAP/microtubule affinity-regulating kinase 1; PAR1 homolog c; Par-1c; Par1c; MARK2; EMK1; Serine/threonine-protein kinase MARK2; ELKL motif kinase 1; EMK-1; MAP/microtubule affin  Immunogen Species Homo sapiens (Human)  Target Names MARK1/MARK2/MARK3/MARK4	Conjugate	Non-conjugated
chromatography using epitope-specific immunogen.  Isotype  IgG  MARK1; KIAA1477; MARK; Serine/threonine-protein kinase MARK1; MAP/microtubule affinity-regulating kinase 1; PAR1 homolog c; Par-1c; Par1c; MARK2; EMK1; Serine/threonine-protein kinase MARK2; ELKL motif kinase 1; EMK-1; MAP/microtubule affin  Immunogen Species  Homo sapiens (Human)  Target Names  MARK1/MARK2/MARK3/MARK4	Storage Buffer	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Alias  MARK1; KIAA1477; MARK; Serine/threonine-protein kinase MARK1;  MAP/microtubule affinity-regulating kinase 1; PAR1 homolog c; Par-1c; Par1c;  MARK2; EMK1; Serine/threonine-protein kinase MARK2; ELKL motif kinase 1;  EMK-1; MAP/microtubule affin  Immunogen Species  Homo sapiens (Human)  Target Names  MARK1/MARK2/MARK3/MARK4	Purification Method	
MAP/microtubule affinity-regulating kinase 1; PAR1 homolog c; Par-1c; Par1c; MARK2; EMK1; Serine/threonine-protein kinase MARK2; ELKL motif kinase 1; EMK-1; MAP/microtubule affin  Immunogen Species Homo sapiens (Human)  Target Names MARK1/MARK2/MARK3/MARK4  Image	Isotype	IgG
Target Names MARK1/MARK2/MARK3/MARK4 Image	Alias	MAP/microtubule affinity-regulating kinase 1; PAR1 homolog c; Par-1c; Par1c; MARK2; EMK1; Serine/threonine-protein kinase MARK2; ELKL motif kinase 1;
Image	Immunogen Species	Homo sapiens (Human)
Image Western Plot analysis of COC7 as its visitors	Target Names	MARK1/MARK2/MARK3/MARK4
	Image	Western Diet englisse of COCZ selle weige



Western Blot analysis of COS7 cells using MARK1/2/3/4 Polyclonal Antibody