





## PLCG1 Antibody

Product CodeCSB-PA003818StorageUpon receipt, store at -20°C or -80°C. Avoid repeated freeze.Uniprot No.P19174ImmunogenSynthesized peptide derived from Human PLC γ1 around the non-phosphorylation site of Y771.Raised InRabbitSpecies ReactivityHuman,Mouse,Rat,MonkeyTested ApplicationsWB, IHC, ELISA; WB:1:500-1:2000, IHC:1:100-1:300, ELISA:1:10000
Uniprot No.       P19174         Immunogen       Synthesized peptide derived from Human PLC γ1 around the non-phosphorylation site of Y771.         Raised In       Rabbit         Species Reactivity       Human,Mouse,Rat,Monkey
Immunogen         Synthesized peptide derived from Human PLC γ1 around the non-phosphorylation site of Y771.           Raised In         Rabbit           Species Reactivity         Human,Mouse,Rat,Monkey
phosphorylation site of Y771.  Raised In Rabbit  Species Reactivity Human,Mouse,Rat,Monkey
Species Reactivity Human, Mouse, Rat, Monkey
<b>Tested Applications</b> WB, IHC, ELISA; WB:1:500-1:2000, IHC:1:100-1:300, ELISA:1:10000
Form Liquid
Conjugate Non-conjugated
Storage Buffer Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azid
Purification Method The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Isotype</b> IgG
Alias  PLCG1; PLC1; 1-phosphatidylinositol 4; 5-bisphosphate phosphodiesteras gamma-1; PLC-148; Phosphoinositide phospholipase C-gamma-1; Phospholipase C-II; PLC-II; Phospholipase C-gamma-1; PLC-gamma-1
Immunogen Species Homo sapiens (Human)
Target Names PLCG1
Image Western Blot analysis of HenG2 cells using BLC

(kD) % 170-130-95-72-55Western Blot analysis of HepG2 cells using PLC  $\gamma$ 1 Polyclonal Antibody