

🕜 Tel: +1-301-363-4651 🛛 🖂 Email: cusabio@cusabio.com 🥃 Website: www.cusabio.com

MUC1 Antibody

StorageUpon receipt, store at -20°C or -80°C. Avoid repeated freeze.Uniprot No.P15941ImmunogenSynthesized peptide derived from Human Mucin 1 around the non- phosphorylation site of S1227.Raised InRabbitSpecies ReactivityHumanTested ApplicationsIHC, ELISA; IHC:1:100-1:300, ELISA:1:10000FormLiquidConjugateNon-conjugatedStorage BufferLiquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.Purification MethodThe antibody was affinity-purified from rabbit antiserum by affinity- chromatography using epitope-specific immunogen.IsotypeIgGAliasMUC1; PUM; Mucin-1; MUC-1; Breast carcinoma-associated antigen DF3; Carcinoma-associated mucin; Episialin; H23AG; Krebs von den Lungen-6; KL-6; PEMT; Peanut-reactive urinary mucin; PUM; Polymorphic epithelial mucin; PEM; Tumor-associated epImmunogen SpeciesHomo sapiens (Human)		
Uniprot No.P15941ImmunogenSynthesized peptide derived from Human Mucin 1 around the non-phosphorylation site of S1227.Raised InRabbitSpecies ReactivityHumanTested ApplicationsIHC, ELISA; IHC:1:100-1:300, ELISA:1:10000FormLiquidConjugateNon-conjugatedStorage BufferLiquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.Purification MethodThe antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.IsotypeIgGAliasMUC1; PUM; Mucin-1; MUC-1; Breast carcinoma-associated antigen DF3; Carcinoma-associated mucin; Episialin; H23AG; Krebs von den Lungen-6; KL-6; PEMT; Peanut-reactive urinary mucin; PUM; Polymorphic epithelial mucin; PEM; Tumor-associated epImmunogen SpeciesHomo sapiens (Human)	Product Code	CSB-PA010276
ImmunogenSynthesized peptide derived from Human Mucin 1 around the non- phosphorylation site of S1227.Raised InRabbitSpecies ReactivityHumanTested ApplicationsIHC, ELISA; IHC:1:100-1:300, ELISA:1:10000FormLiquidConjugateNon-conjugatedStorage BufferLiquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.Purification MethodThe antibody was affinity-purified from rabbit antiserum by affinity- chromatography using epitope-specific immunogen.IsotypeIgGAliasMUC1; PUM; Mucin-1; MUC-1; Breast carcinoma-associated antigen DF3; Carcinoma-associated mucin; Episialin; H23AG; Krebs von den Lungen-6; KL-6; PEMT; Peanut-reactive urinary mucin; PUM; Polymorphic epithelial mucin; PEM; Tumor-associated epImmunogen SpeciesHomo sapiens (Human)	Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Phosphorylation site of S1227.Raised InRabbitSpecies ReactivityHumanTested ApplicationsIHC, ELISA; IHC:1:100-1:300, ELISA:1:10000FormLiquidConjugateNon-conjugatedStorage BufferLiquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.Purification MethodThe antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.IsotypeIgGAliasMUC1; PUM; Mucin-1; MUC-1; Breast carcinoma-associated antigen DF3; Carcinoma-associated mucin; Episialin; H23AG; Krebs von den Lungen-6; KL-6; PEMT; Peanut-reactive urinary mucin; PUM; Polymorphic epithelial mucin; PEM; Tumor-associated epImmunogen SpeciesHomo sapiens (Human)	Uniprot No.	P15941
Species Reactivity Human Tested Applications IHC, ELISA; IHC:1:100-1:300, ELISA:1:10000 Form Liquid Conjugate Non-conjugated Storage Buffer Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Purification Method The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Isotype IgG Alias MUC1; PUM; Mucin-1; MUC-1; Breast carcinoma-associated antigen DF3; Carcinoma-associated mucin; Episialin; H23AG; Krebs von den Lungen-6; KL-6; PEMT; Peanut-reactive urinary mucin; PUM; Polymorphic epithelial mucin; PEM; Tumor-associated ep Immunogen Species Homo sapiens (Human)	Immunogen	
Tested ApplicationsIHC, ELISA; IHC:1:100-1:300, ELISA:1:10000FormLiquidConjugateNon-conjugatedStorage BufferLiquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.Purification MethodThe antibody was affinity-purified from rabbit antiserum by affinity- chromatography using epitope-specific immunogen.IsotypeIgGAliasMUC1; PUM; Mucin-1; MUC-1; Breast carcinoma-associated antigen DF3; Carcinoma-associated mucin; Episialin; H23AG; Krebs von den Lungen-6; KL-6; PEMT; Peanut-reactive urinary mucin; PUM; Polymorphic epithelial mucin; PEM; Tumor-associated epImmunogen SpeciesHomo sapiens (Human)	Raised In	Rabbit
FormLiquidConjugateNon-conjugatedStorage BufferLiquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.Purification MethodThe antibody was affinity-purified from rabbit antiserum by affinity- chromatography using epitope-specific immunogen.IsotypeIgGAliasMUC1; PUM; Mucin-1; MUC-1; Breast carcinoma-associated antigen DF3; Carcinoma-associated mucin; Episialin; H23AG; Krebs von den Lungen-6; KL-6; PEMT; Peanut-reactive urinary mucin; PUM; Polymorphic epithelial mucin; PEM; Tumor-associated epImmunogen SpeciesHomo sapiens (Human)	Species Reactivity	Human
ConjugateNon-conjugatedStorage BufferLiquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.Purification MethodThe antibody was affinity-purified from rabbit antiserum by affinity- chromatography using epitope-specific immunogen.IsotypeIgGAliasMUC1; PUM; Mucin-1; MUC-1; Breast carcinoma-associated antigen DF3; Carcinoma-associated mucin; Episialin; H23AG; Krebs von den Lungen-6; KL-6; PEMT; Peanut-reactive urinary mucin; PUM; Polymorphic epithelial mucin; PEM; Tumor-associated epImmunogen SpeciesHomo sapiens (Human)	Tested Applications	IHC, ELISA; IHC:1:100-1:300, ELISA:1:10000
Storage BufferLiquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.Purification MethodThe antibody was affinity-purified from rabbit antiserum by affinity- chromatography using epitope-specific immunogen.IsotypeIgGAliasMUC1; PUM; Mucin-1; MUC-1; Breast carcinoma-associated antigen DF3; Carcinoma-associated mucin; Episialin; H23AG; Krebs von den Lungen-6; KL-6; PEMT; Peanut-reactive urinary mucin; PUM; Polymorphic epithelial mucin; PEM; Tumor-associated epImmunogen SpeciesHomo sapiens (Human)	Form	Liquid
Purification Method The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Isotype IgG Alias MUC1; PUM; Mucin-1; MUC-1; Breast carcinoma-associated antigen DF3; Carcinoma-associated mucin; Episialin; H23AG; Krebs von den Lungen-6; KL-6; PEMT; Peanut-reactive urinary mucin; PUM; Polymorphic epithelial mucin; PEM; Tumor-associated ep Immunogen Species Homo sapiens (Human)	Conjugate	Non-conjugated
Isotype IgG Alias MUC1; PUM; Mucin-1; MUC-1; Breast carcinoma-associated antigen DF3; Carcinoma-associated mucin; Episialin; H23AG; Krebs von den Lungen-6; KL-6; PEMT; Peanut-reactive urinary mucin; PUM; Polymorphic epithelial mucin; PEM; Tumor-associated ep Immunogen Species Homo sapiens (Human)	Storage Buffer	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Alias MUC1; PUM; Mucin-1; MUC-1; Breast carcinoma-associated antigen DF3; Carcinoma-associated mucin; Episialin; H23AG; Krebs von den Lungen-6; KL-6; PEMT; Peanut-reactive urinary mucin; PUM; Polymorphic epithelial mucin; PEM; Tumor-associated ep Immunogen Species Homo sapiens (Human)	Purification Method	
Carcinoma-associated mucin; Episialin; H23AG; Krebs von den Lungen-6; KL-6; PEMT; Peanut-reactive urinary mucin; PUM; Polymorphic epithelial mucin; PEM; Tumor-associated ep Immunogen Species Homo sapiens (Human)	Isotype	lgG
	Alias	Carcinoma-associated mucin; Episialin; H23AG; Krebs von den Lungen-6; KL-6; PEMT; Peanut-reactive urinary mucin; PUM; Polymorphic epithelial mucin;
Target Names MUC1	Immunogen Species	Homo sapiens (Human)
	Target Names	MUC1

1