







## TGM4 Antibody

| <b>Product Code</b>                        | CSB-PA038312   |
|--|--|
| Storage                                    | Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.                                |
| Uniprot No.                                | P49221   |
| Immunogen                                  | Fusion protein of Human TGM4   |
| Raised In                                  | Rabbit   |
| Species Reactivity                         | Human  |
| <b>Tested Applications</b>                 | ELISA,WB,IHC;ELISA:1:2000-1:5000,WB:1:500-1:2000,IHC:1:25-1:100                              |
| Form                                       | Liquid   |
|  | · d · ·  |
| Conjugate                                  | Non-conjugated   |
| Conjugate Storage Buffer                   | •  |
|  | Non-conjugated   |
| Storage Buffer                             | Non-conjugated -20°C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol                                    |
| Storage Buffer Purification Method         | Non-conjugated -20°C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol Antigen affinity purification      |
| Storage Buffer Purification Method Isotype | Non-conjugated -20°C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol Antigen affinity purification  IgG |





The image on the left is immunohistochemistry of paraffin-embedded Human prostate cancer tissue using CSB-PA038312(TGM4 Antibody) at dilution 1/20, on the right is treated with fusion protein. (Original magnification: ×200)



The image on the left is immunohistochemistry of paraffin-embedded Human esophagus cancer tissue using CSB-PA038312(TGM4 Antibody) at dilution 1/20, on the right is treated with fusion protein. (Original magnification: ×200)




## **CUSABIO TECHNOLOGY LLC**

🕜 Tel: +1-301-363-4651 🛛 Email: cusabio@cusabio.com 🌔 Website: www.cusabio.com 🍵







Gel: 6%SDS-PAGE, Lysate: 40  $\mu$ g, Lane 1-5: Hela, Incap, skov3, 293T and TM4 cell, Primary antibody: CSB-PA038312(TGM4 Antibody) at dilution 1/400 dilution, Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution, Exposure time: 10 seconds