


🕜 Tel: +1-301-363-4651 🛛 🖂 Email: cusabio@cusabio.com 🥥 Website: www.cusabio.com 🧉

DDX50 Antibody

StorageUpon receipt, store at -20°C or -80°C. Avoid repeated freeze.Jniprot No.Q9BQ39mmunogenRecombinant Human ATP-dependent RNA helicase DDX50 protein (1-210AA)Raised InRabbitSpecies ReactivityHumanFested ApplicationsELISA, WB, IHC; Recommended dilution: WB:1:500-1:2000, IHC:1:20-1:200FormLiquidConjugateNon-conjugatedStorage BufferPreservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4Purification Method>95%, Protein G purifiedstorageIgGClonalityPolyclonalMiasATP-dependent RNA helicase DDX50 (EC 3.6.4.13) (DEAD box protein 50) (Gu-beta) (Nucleolar protein Gu2), DDX50munogen SpeciesHomo sapiens (Human)Research AreaEpigenetics and Nuclear SignalingTarget NamesDDX50		
Jniprot No.Q9BQ39ImmunogenRecombinant Human ATP-dependent RNA helicase DDX50 protein (1-210AA)Raised InRabbitSpecies ReactivityHumanTested ApplicationsELISA, WB, IHC; Recommended dilution: WB:1:500-1:2000, IHC:1:20-1:200FormLiquidConjugateNon-conjugatedStorage BufferPreservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4Purification Method>95%, Protein G purifiedStorageIgGClonalityPolyclonalNiasATP-dependent RNA helicase DDX50 (EC 3.6.4.13) (DEAD box protein 50) (Gu-beta) (Nucleolar protein Gu2), DDX50Immunogen SpeciesHomo sapiens (Human)Research AreaEpigenetics and Nuclear SignalingmageDDX50	Product Code	CSB-PA861080LA01HU
mmunogenRecombinant Human ATP-dependent RNA helicase DDX50 protein (1-210AA)Raised InRabbitSpecies ReactivityHumanTested ApplicationsELISA, WB, IHC; Recommended dilution: WB:1:500-1:2000, IHC:1:20-1:200FormLiquidConjugateNon-conjugatedStorage BufferPreservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4Purification Method>95%, Protein G purifiedstorageIgGClonalityPolyclonalMiasATP-dependent RNA helicase DDX50 (EC 3.6.4.13) (DEAD box protein 50) (Gu-beta) (Nucleolar protein Gu2), DDX50mmunogen SpeciesHomo sapiens (Human)Research AreaEpigenetics and Nuclear Signalingmage	Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Raised InRabbitRabbitHumanSpecies ReactivityHumanTested ApplicationsELISA, WB, IHC; Recommended dilution: WB:1:500-1:2000, IHC:1:20-1:200FormLiquidConjugateNon-conjugatedStorage BufferPreservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4Purification Method>95%, Protein G purifiedstotypeIgGClonalityPolyclonalNiasATP-dependent RNA helicase DDX50 (EC 3.6.4.13) (DEAD box protein 50) (Gu-beta) (Nucleolar protein Gu2), DDX50mmunogen SpeciesHomo sapiens (Human)Research AreaEpigenetics and Nuclear SignalingmageDDX50	Uniprot No.	Q9BQ39
Species Reactivity Human Tested Applications ELISA, WB, IHC; Recommended dilution: WB:1:500-1:2000, IHC:1:20-1:200 Form Liquid Conjugate Non-conjugated Storage Buffer Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 Purification Method >95%, Protein G purified Storage IgG Clonality Polyclonal Alias ATP-dependent RNA helicase DDX50 (EC 3.6.4.13) (DEAD box protein 50) (Gu-beta) (Nucleolar protein Gu2), DDX50 mmunogen Species Homo sapiens (Human) Research Area Epigenetics and Nuclear Signaling mage DDX50	Immunogen	Recombinant Human ATP-dependent RNA helicase DDX50 protein (1-210AA)
Tested ApplicationsELISA, WB, IHC; Recommended dilution: WB:1:500-1:2000, IHC:1:20-1:200FormLiquidConjugateNon-conjugatedStorage BufferPreservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4Purification Method>95%, Protein G purifiedsotypeIgGClonalityPolyclonalAliasATP-dependent RNA helicase DDX50 (EC 3.6.4.13) (DEAD box protein 50) (Gu-beta) (Nucleolar protein Gu2), DDX50mmunogen SpeciesHomo sapiens (Human)Research AreaEpigenetics and Nuclear SignalingmageDDX50	Raised In	Rabbit
FormLiquidConjugateNon-conjugatedStorage BufferPreservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4Purification Method>95%, Protein G purifiedSotypeIgGClonalityPolyclonalNiasATP-dependent RNA helicase DDX50 (EC 3.6.4.13) (DEAD box protein 50) (Gu-beta) (Nucleolar protein Gu2), DDX50mmunogen SpeciesHomo sapiens (Human)Research AreaEpigenetics and Nuclear SignalingmageImage	Species Reactivity	Human
ConjugateNon-conjugatedStorage BufferPreservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4Purification Method>95%, Protein G purifiedsotypeIgGClonalityPolyclonalNiasATP-dependent RNA helicase DDX50 (EC 3.6.4.13) (DEAD box protein 50) (Gu-beta) (Nucleolar protein Gu2), DDX50mmunogen SpeciesHomo sapiens (Human)Research AreaEpigenetics and Nuclear SignalingTarget NamesDDX50	Tested Applications	ELISA, WB, IHC; Recommended dilution: WB:1:500-1:2000, IHC:1:20-1:200
Storage BufferPreservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4Purification Method>95%, Protein G purifiedSotypeIgGClonalityPolyclonalNiasATP-dependent RNA helicase DDX50 (EC 3.6.4.13) (DEAD box protein 50) (Gu-beta) (Nucleolar protein Gu2), DDX50mmunogen SpeciesHomo sapiens (Human)Research AreaEpigenetics and Nuclear SignalingTarget NamesDDX50	Form	Liquid
Constituents: 50% Glycerol, 0.01M PBS, PH 7.4Purification Method>95%, Protein G purifiedsotypeIgGClonalityPolyclonalNiasATP-dependent RNA helicase DDX50 (EC 3.6.4.13) (DEAD box protein 50) (Gu-beta) (Nucleolar protein Gu2), DDX50mmunogen SpeciesHomo sapiens (Human)Research AreaEpigenetics and Nuclear SignalingTarget NamesDDX50	Conjugate	Non-conjugated
sotypeIgGClonalityPolyclonalAliasATP-dependent RNA helicase DDX50 (EC 3.6.4.13) (DEAD box protein 50) (Gu-beta) (Nucleolar protein Gu2), DDX50mmunogen SpeciesHomo sapiens (Human)Research AreaEpigenetics and Nuclear SignalingTarget NamesDDX50	Storage Buffer	
ClonalityPolyclonalAliasATP-dependent RNA helicase DDX50 (EC 3.6.4.13) (DEAD box protein 50) (Gu-beta) (Nucleolar protein Gu2), DDX50mmunogen SpeciesHomo sapiens (Human)Research AreaEpigenetics and Nuclear SignalingTarget NamesDDX50	Purification Method	>95%, Protein G purified
AliasATP-dependent RNA helicase DDX50 (EC 3.6.4.13) (DEAD box protein 50) (Gu-beta) (Nucleolar protein Gu2), DDX50mmunogen SpeciesHomo sapiens (Human)Research AreaEpigenetics and Nuclear SignalingTarget NamesDDX50	Isotype	lgG
(Gu-beta) (Nucleolar protein Gu2), DDX50 mmunogen Species Homo sapiens (Human) Research Area Epigenetics and Nuclear Signaling Target Names DDX50	Clonality	Polyclonal
Research Area Epigenetics and Nuclear Signaling Farget Names DDX50 mage	Alias	
Target Names DDX50	Immunogen Species	Homo sapiens (Human)
mage	Research Area	Epigenetics and Nuclear Signaling
mage Western blot	Target Names	DDX50
	Image	Western blot




1



CUSABIO TECHNOLOGY LLC

🕜 Tel: +1-301-363-4651 🛛 🖂 Email: cusabio@cusabio.com 🥥 Website: www.cusabio.com 🧉





Immunohistochemistry of paraffin-embedded human thyroid tissue using CSB-PA861080LA01HU at dilution of 1:100