




FAIM Antibody

| | |
|----------------------------|--|
| Product Code | CSB-PA889122LA01HU |
| Storage | Upon receipt, store at -20°C or -80°C. Avoid repeated freeze. |
| Uniprot No. | Q9NVQ4 |
| Immunogen | Recombinant Human Fas apoptotic inhibitory molecule 1 protein (1-179AA) |
| Raised In | Rabbit |
| Species Reactivity | Human, Rat |
| Tested Applications | ELISA, WB, IHC, IF; Recommended dilution: WB:1:500-1:5000, IHC:1:20-1:200, IF:1:50-1:200 |
| Form | Liquid |
| Conjugate | Non-conjugated |
| Storage Buffer | Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 |
| Purification Method | >95%, Protein G purified |
| Isotype | IgG |
| Clonality | Polyclonal |
| Alias | Fas apoptotic inhibitory molecule 1, FAIM, FAIM1 |
| Immunogen Species | Homo sapiens (Human) |
| Research Area | Cell Biology |
| Target Names | FAIM |

Image



Western blot

All lanes: FAIM antibody at 2µg/ml

Lane 1: 293T whole cell lysate

Lane 2: Rat gonad tissue


Lane 3: Jurkat whole cell lysate

Secondary


Goat polyclonal to rabbit IgG at 1/10000 dilution

Predicted band size: 21, 25, 23 kDa

Observed band size: 21 kDa



Immunohistochemistry of paraffin-embedded human breast cancer using CSB-PA889122LA01HU at dilution of 1:100



Immunofluorescence staining of HepG2 cells with CSB-PA889122LA01HU at 1:133, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).

Usage

For Research Use Only. Not for use in diagnostic or therapeutic procedures.