



🕧 Tel: +1-301-363-4651 🛛 🖂 Email: cusabio@cusabio.com 📀 Website: www.cusabio.com 🌘

FXYD7 Antibody

diameter(47-80AA)Raised InRabbitSpecies ReactivityHumanTested ApplicationsELISA, IHC, IF; Recommended dilution: IHC:1:20-1:200, IF:1:50-1:200FormLiquidConjugateNon-conjugatedStorage BufferPreservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4Purification Method>95%, Protein G purifiedIsotypeIgGClonalityPolyclonalAliasFXYD domain-containing ion transport regulator 7, FXYD7Immunogen SpeciesHomo sapiens (Human)Research AreaSignal TransductionImage		
Uniprot No.P58549ImmunogenRecombinant Human FXYD domain-containing ion transport regulator 7 prote (47-80AA)Raised InRabbitSpecies ReactivityHumanTested ApplicationsELISA, IHC, IF; Recommended dilution: IHC:1:20-1:200, IF:1:50-1:200FormLiquidConjugateNon-conjugatedStorage BufferPreservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4Purification Method>95%, Protein G purifiedIsotypeIgGClonalityPolyclonalAliasFXYD domain-containing ion transport regulator 7, FXYD7Immunogen SpeciesHomo sapiens (Human)Research AreaSignal TransductionTarget NamesFXYD7	Product Code	CSB-PA009097LA01HU
ImmunogenRecombinant Human FXYD domain-containing ion transport regulator 7 protect (47-80AA)Raised InRabbitSpecies ReactivityHumanTested ApplicationsELISA, IHC, IF; Recommended dilution: IHC:1:20-1:200, IF:1:50-1:200FormLiquidConjugateNon-conjugatedStorage BufferPreservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4Purification Method>95%, Protein G purifiedIsotypeIgGClonalityPolyclonalAliasFXYD domain-containing ion transport regulator 7, FXYD7Immunogen SpeciesHomo sapiens (Human)Research AreaSignal TransductionTarget NamesFXYD7	Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
aised InRabbitSpecies ReactivityHumanTested ApplicationsELISA, IHC, IF; Recommended dilution: IHC:1:20-1:200, IF:1:50-1:200FormLiquidConjugateNon-conjugatedStorage BufferPreservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4Purification Method>95%, Protein G purifiedIsotypeIgGClonalityPolyclonalAliasFXYD domain-containing ion transport regulator 7, FXYD7Immunogen SpeciesHomo sapiens (Human)Research AreaSignal TransductionImage	Uniprot No.	P58549
Species ReactivityHumanTested ApplicationsELISA, IHC, IF; Recommended dilution: IHC:1:20-1:200, IF:1:50-1:200FormLiquidConjugateNon-conjugatedStorage BufferPreservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4Purification Method>95%, Protein G purifiedIsotypeIgGClonalityPolyclonalAliasFXYD domain-containing ion transport regulator 7, FXYD7Immunogen SpeciesHomo sapiens (Human)Research AreaSignal TransductionTarget NamesFXYD7	Immunogen	Recombinant Human FXYD domain-containing ion transport regulator 7 protein (47-80AA)
Tested ApplicationsELISA, IHC, IF; Recommended dilution: IHC:1:20-1:200, IF:1:50-1:200FormLiquidConjugateNon-conjugatedStorage BufferPreservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4Purification Method>95%, Protein G purifiedIsotypeIgGClonalityPolyclonalAliasFXYD domain-containing ion transport regulator 7, FXYD7Immunogen SpeciesHomo sapiens (Human)Research AreaSignal TransductionTarget NamesFXYD7	Raised In	Rabbit
FormLiquidConjugateNon-conjugatedStorage BufferPreservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4Purification Method>95%, Protein G purifiedIsotypeIgGClonalityPolyclonalAliasFXYD domain-containing ion transport regulator 7, FXYD7Immunogen SpeciesHomo sapiens (Human)Research AreaSignal TransductionImageEXYD7	Species Reactivity	Human
ConjugateNon-conjugatedStorage BufferPreservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4Purification Method>95%, Protein G purifiedIsotypeIgGClonalityPolyclonalAliasFXYD domain-containing ion transport regulator 7, FXYD7Immunogen SpeciesHomo sapiens (Human)Research AreaSignal TransductionTarget NamesFXYD7	Tested Applications	ELISA, IHC, IF; Recommended dilution: IHC:1:20-1:200, IF:1:50-1:200
Storage BufferPreservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4Purification Method>95%, Protein G purifiedIsotypeIgGClonalityPolyclonalAliasFXYD domain-containing ion transport regulator 7, FXYD7Immunogen SpeciesHomo sapiens (Human)Research AreaSignal TransductionTarget NamesFXYD7	Form	Liquid
Constituents: 50% Glycerol, 0.01M PBS, PH 7.4Purification Method>95%, Protein G purifiedIsotypeIgGClonalityPolyclonalAliasFXYD domain-containing ion transport regulator 7, FXYD7Immunogen SpeciesHomo sapiens (Human)Research AreaSignal TransductionTarget NamesFXYD7	Conjugate	Non-conjugated
IsotypeIgGClonalityPolyclonalAliasFXYD domain-containing ion transport regulator 7, FXYD7Immunogen SpeciesHomo sapiens (Human)Research AreaSignal TransductionTarget NamesFXYD7	Storage Buffer	
Clonality Polyclonal Alias FXYD domain-containing ion transport regulator 7, FXYD7 Immunogen Species Homo sapiens (Human) Research Area Signal Transduction Target Names FXYD7	Purification Method	>95%, Protein G purified
Alias FXYD domain-containing ion transport regulator 7, FXYD7 Immunogen Species Homo sapiens (Human) Research Area Signal Transduction Target Names FXYD7	Isotype	lgG
Immunogen Species Homo sapiens (Human) Research Area Signal Transduction Target Names FXYD7 Image	Clonality	Polyclonal
Research Area Signal Transduction Target Names FXYD7 Image	Alias	FXYD domain-containing ion transport regulator 7, FXYD7
Target Names FXYD7	Immunogen Species	Homo sapiens (Human)
	Research Area	Signal Transduction
Image	Target Names	FXYD7
	Image	Immunohistochemistry of paraffin-embedded



Immunohistochemistry of paraffin-embedded human heart tissue using CSB-PA009097LA01HU at dilution of 1:100



Immunofluorescent analysis of U251 cells using CSB-PA009097LA01HU at dilution of 1:100 and Alexa Fluor 488-congugated AffiniPure Goat Anti-Rabbit IgG(H+L)

1