




# PORCN Antibody

<b>Product Code</b>	CSB-PA887958LA01HU
<b>Storage</b>	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
<b>Uniprot No.</b>	Q9H237
<b>Immunogen</b>	Recombinant Human Protein-serine O-palmitoleoyltransferase porcupine protein (269-332AA)
<b>Raised In</b>	Rabbit
<b>Species Reactivity</b>	Human, Mouse
<b>Tested Applications</b>	ELISA, WB, IHC, IF; Recommended dilution: WB:1:500-1:2000, IHC:1:20-1:200, IF:1:50-1:200
<b>Form</b>	Liquid
<b>Conjugate</b>	Non-conjugated
<b>Storage Buffer</b>	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4
<b>Purification Method</b>	>95%, Protein G purified
<b>Isotype</b>	IgG
<b>Clonality</b>	Polyclonal
<b>Alias</b>	Protein-serine O-palmitoleoyltransferase porcupine (EC 2.3.1.-) (Protein MG61), PORCN, MG61 PORC PPN
<b>Immunogen Species</b>	Homo sapiens (Human)
<b>Research Area</b>	Tags & Cell Markers
<b>Target Names</b>	PORCN

## Image



### Western blot


All lanes: PORCN antibody at 3μg/ml + Mouse thymus tissue

### Secondary


Goat polyclonal to rabbit IgG at 1/10000 dilution

Predicted band size: 53, 52, 43 kDa

Observed band size: 53 kDa



Immunohistochemistry of paraffin-embedded human testis tissue using CSB-PA887958LA01HU at dilution of 1:100



Immunofluorescence staining of A549 cells with CSB-PA887958LA01HU at 1:100, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).