




# TTYH1 Antibody

<b>Product Code</b>	CSB-PA867139LA01HU
<b>Storage</b>	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
<b>Uniprot No.</b>	Q9H313
<b>Immunogen</b>	Synthesized peptide derived from human TTYH1
<b>Raised In</b>	Rabbit
<b>Species Reactivity</b>	Human,mouse
<b>Tested Applications</b>	ELISA, WB, IF; Recommended dilution: WB:1:500-1:1000, IF:1:20-1:50
<b>Form</b>	Liquid
<b>Conjugate</b>	Non-conjugated
<b>Storage Buffer</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Purification Method</b>	Antigen affinity purification
<b>Isotype</b>	IgG
<b>Clonality</b>	Polyclonal
<b>Product Type</b>	Polyclonal Antibody
<b>Immunogen Species</b>	Homo sapiens (Human)
<b>Research Area</b>	Others
<b>Target Names</b>	TTYH1

## Image



### Western Blot

Positive WB detected in: Raji whole cell lysate(20µg), HeLa whole cell lysate(20µg), HepG2 whole cell lysate(20µg), Mouse Brain tissue lysate(20µg)

All lanes: TTYH1 antibody at 1:1000


Secondary

Goat polyclonal to rabbit IgG at 1/20000 dilution


Predicted band size: 50, 27,36 kDa

Observed band size: 50 kDa

Exposure time?120s



Immunofluorescence staining of HeLa cell with CSB-PA867139LA01HU at 1:30, counter-stained with DAPI. The cells were fixed in 4% formaldehyde and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4C. The secondary antibody was Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Immunofluorescence staining of HeLa cell with 5% goat serum, counter-stained with DAPI. The cells were fixed in 4% formaldehyde and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4C. The secondary antibody was Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).

---

**Usage**

For Research Use Only. Not for use in diagnostic or therapeutic procedures.