




# UXS1 Antibody

<b>Product Code</b>	CSB-PA822754LA01HU
<b>Storage</b>	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
<b>Uniprot No.</b>	Q8NBZ7
<b>Immunogen</b>	Recombinant Human UDP-glucuronic acid decarboxylase 1 protein (201-420AA)
<b>Raised In</b>	Rabbit
<b>Species Reactivity</b>	Human
<b>Tested Applications</b>	ELISA, WB, IHC; Recommended dilution: WB:1:500-1:2000, IHC:1:1:50-1:200
<b>Form</b>	Liquid
<b>Conjugate</b>	Non-conjugated
<b>Storage Buffer</b>	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4
<b>Purification Method</b>	Antigen affinity purification
<b>Isotype</b>	IgG
<b>Clonality</b>	Polyclonal
<b>Product Type</b>	Polyclonal Antibody
<b>Immunogen Species</b>	Homo sapiens (Human)
<b>Research Area</b>	Others
<b>Target Names</b>	UXS1

## Image



### Western Blot

Positive WB detected in: JK whole cell lysate(20µg), HepG2 whole cell lysate(20µg), PC-3 whole cell lysate(20µg), A375 whole cell lysate(20µg)

All lanes: UXS1 antibody at 1:1000


Secondary

Goat polyclonal to rabbit IgG at 1/20000 dilution


Predicted band size: 48,49,29 kDa

Observed band size: 70-100 kDa

Exposure time: 120s



IHC image of CSB-PA822754LA01HU diluted at 1:50 and staining in paraffin-embedded human Lung tissue performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a Goat anti-rabbit polymer IgG labeled by HRP and visualized using 0.05% DAB. Secondary antibody only control: uses 1% BSA instead of primary antibody



IHC image of CSB-PA822754LA01HU diluted at 1:50 and staining in paraffin-embedded human Stomach tissue performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a Goat anti-rabbit polymer IgG labeled by HRP and visualized using 0.05% DAB. Secondary antibody only control: uses 1% BSA instead of primary antibody

**Usage**

For Research Use Only. Not for use in diagnostic or therapeutic procedures.