




FNTB Antibody

Product Code	CSB-PA008780LA01HU
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Uniprot No.	P49356
Immunogen	Recombinant Human Protein farnesyltransferase subunit beta protein (1-300AA)
Raised In	Rabbit
Species Reactivity	Human
Tested Applications	ELISA, WB, IHC, IF, IP; Recommended dilution: WB:1:500-1:5000, IHC:1:20-1:200, IF:1:50-1:200, IP:1:200-1:2000
Form	Liquid
Conjugate	Non-conjugated
Storage Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4
Purification Method	>95%, Protein G purified
Isotype	IgG
Clonality	Polyclonal
Alias	Protein farnesyltransferase subunit beta (FTase-beta) (EC 2.5.1.58) (CAAX farnesyltransferase subunit beta) (Ras proteins prenyltransferase subunit beta), FNTB
Immunogen Species	Homo sapiens (Human)
Research Area	Signal Transduction
Target Names	FNTB

Image



Western blot

All lanes: FNTB antibody at 6µg/ml

Lane 1: HeLa whole cell lysate

Lane 2: CEM whole cell lysate

Lane 3: HepG2 whole cell lysate

Lane 4: SH-SY5Y whole cell lysate


Lane 5: A549 whole cell lysate

Secondary


Goat polyclonal to rabbit IgG at 1/10000 dilution

Predicted band size: 49, 44 kDa


Observed band size: 49 kDa



Immunohistochemistry of paraffin-embedded human tonsil tissue using CSB-PA008780LA01HU at dilution of 1:100



Immunofluorescent analysis of HepG2 cells using CSB-PA008780LA01HU at dilution of 1:100 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L)



Immunoprecipitating FNTB in HepG2 whole cell lysate
 Lane 1: Rabbit control IgG (1μg) instead of CSB-PA008780LA01HU in HepG2 whole cell lysate. For western blotting, a HRP-conjugated Protein G antibody was used as the secondary antibody (1/2000)
 Lane 2: CSB-PA008780LA01HU (8μg) + HepG2 whole cell lysate (500μg)
 Lane 3: HepG2 whole cell lysate (10μg)