




TCTA Antibody

| | |
|----------------------------|--|
| Product Code | CSB-PA023331LA01HU |
| Storage | Upon receipt, store at -20°C or -80°C. Avoid repeated freeze. |
| Uniprot No. | P57738 |
| Immunogen | Recombinant Human T-cell leukemia translocation-altered gene protein (62-103AA) |
| Raised In | Rabbit |
| Species Reactivity | Human, Mouse |
| Tested Applications | ELISA, WB, IHC; Recommended dilution: WB:1:1000-1:5000, IHC:1:20-1:200 |
| Form | Liquid |
| Conjugate | Non-conjugated |
| Storage Buffer | Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 |
| Purification Method | Antigen Affinity Purified |
| Isotype | IgG |
| Clonality | Polyclonal |
| Alias | T-cell leukemia translocation-altered gene protein (T-cell leukemia translocation-associated gene protein), TCTA |
| Immunogen Species | Homo sapiens (Human) |
| Research Area | Signal Transduction |
| Target Names | TCTA |

Image



Western blot

All lanes: TCTA antibody at 1.6µg/ml

Lane 1: Mouse liver tissue

Lane 2: Mouse gonadal tissue

Lane 3: Mouse muscle tissue

Lane 4: HL60 whole cell lysate

Lane 5: U251 whole cell lysate


Lane 6: MCF-7 whole cell lysate

Secondary


Goat polyclonal to rabbit IgG at 1/10000 dilution

Predicted band size: 12 kDa

Observed band size: 12 kDa



Immunohistochemistry of paraffin-embedded human adrenal gland tissue using CSB-PA023331LA01HU at dilution of 1:100



Immunohistochemistry of paraffin-embedded human colon cancer using CSB-PA023331LA01HU at dilution of 1:100

Usage

For Research Use Only. Not for use in diagnostic or therapeutic procedures.