





# ZDHHC12 Antibody

<b>Product Code</b>	CSB-PA822219LA01HU
<b>Storage</b>	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
<b>Uniprot No.</b>	Q96GR4
<b>Immunogen</b>	Recombinant Human Probable palmitoyltransferase ZDHHC12 protein(65-140aa)
<b>Raised In</b>	Rabbit
<b>Species Reactivity</b>	Human, Mouse, Rat
<b>Tested Applications</b>	ELISA, WB,IHC; Recommended dilution: WB:1:500-1:2000,IHC:1:20-1:100
<b>Form</b>	Liquid
<b>Conjugate</b>	Non-conjugated
<b>Storage Buffer</b>	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4
<b>Purification Method</b>	Antigen Affinity Purified
<b>Isotype</b>	IgG
<b>Clonality</b>	Polyclonal
<b>Product Type</b>	Polyclonal Antibody
<b>Immunogen Species</b>	Homo sapiens (Human)
<b>Research Area</b>	Neuroscience
<b>Target Names</b>	ZDHHC12

## Image



IHC image of CSB-PA822219LA01HU diluted at 1:50 and staining in paraffin-embedded human kidney tissue performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a Goat anti-rabbit polymer IgG labeled by HRP and visualized using 0.05% DAB.



## Western Blot

Positive WB detected in: MCF7 whole cell lysate(30µg), COLO205 whole cell lysate(30µg), HEK293 whole cell lysate(30µg), Jurkat whole cell lysate(30µg), U251 whole cell lysate(30µg), Mouse brain tissue lysate(30µg), Rat brain tissue lysate(30µg)  
All lanes: ZDHHC12 antibody at 1:1000  
Secondary  
Goat polyclonal to rabbit IgG at 1/40000 dilution



Predicted band size: 31, 25, 37 kDa  
Observed band size: 37 kDa  
Exposure time: 60s

---

**Usage**

For Research Use Only. Not for use in diagnostic or therapeutic procedures.