


🕜 Tel: +1-301-363-4651 🛛 🖂 Email: cusabio@cusabio.com 🛛 🥑 Website: www.cusabio.com 🌘

FATE1 Antibody

| Product Code | CSB-PA839274LA01HU |
|---------------------|---|
| Storage | Upon receipt, store at -20°C or -80°C. Avoid repeated freeze. |
| Uniprot No. | Q969F0 |
| Immunogen | Recombinant Human Fetal and adult testis-expressed transcript protein (1-162AA) |
| Raised In | Rabbit |
| Species Reactivity | Human |
| Tested Applications | ELISA, IHC, IF; Recommended dilution: IHC:1:200-1:500, IF:1:50-1:500 |
| Form | Liquid |
| Conjugate | Non-conjugated |
| Storage Buffer | Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 |
| Purification Method | >95%, Protein G purified |
| Isotype | IgG |
| Clonality | Polyclonal |
| Alias | Fetal and adult testis-expressed transcript protein (Cancer/testis antigen 43) (CT43) (Tumor antigen BJ-HCC-2), FATE1, FATE |
| Immunogen Species | Homo sapiens (Human) |
| Research Area | Cancer |
| Target Names | FATE1 |
| | |

Image




IHC image of CSB-PA839274LA01HU diluted at 1:400 and staining in paraffin-embedded human testis tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.

1




CUSABIO TECHNOLOGY LLC

🕜 Tel: +1-301-363-4651 🛛 🖂 Email: cusabio@cusabio.com 🛛 🥑 Website: www.cusabio.com 🍙



IHC image of CSB-PA839274LA01HU diluted at 1:400 and staining in paraffin-embedded human ovarian cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



Immunofluorescent analysis of A549 cells using CSB-PA839274LA01HU at dilution of 1:100 and Alexa Fluor 488-congugated AffiniPure Goat Anti-Rabbit IgG(H+L)